



AD-A142 598

DELINEATION OF WETLAND BOUNDARIES USING VEGETATION
WITHIN THE ALTAMHA RIVER BASIN OF GEORGIA(U)
ENVIROSPHERE CO ATLANTA GA B F VAUGHN ET AL APR 84
UNCLASSIFIED WES/TR/Y-84-1 DACW39-78-C-0092 F/G 8/8

1/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A



US Army Corps
of Engineers

AD-A142 590



WETLANDS RESEARCH PROGRAM

TECHNICAL REPORT Y-84-1

DELINEATION OF WETLAND BOUNDARIES USING VEGETATION WITHIN THE ALTAMAHA RIVER BASIN OF GEORGIA

by

Boyd F. Vaughan, Jr., Robert J. Cooper,
Joel H. Braswell, Robin Hart

Envirosphere Company
145 Technology Park
Atlanta, Ga. 30092



April 1984

Final Report

Approved For Public Release. Distribution Unlimited

DTIC
JUN 28 84

DTIC FILE COPY

Prepared for Office, Chief of Engineers, U. S. Army
Washington, D. C. 20314

Under Contract No. DACW39-78-C-0092

Monitored by Environmental Laboratory
U. S. Army Engineer Waterways Experiment Station
P. O. Box 631, Vicksburg, Miss. 39180

84 06 27 080

Destroy this report when no longer needed. Do not
return it to the originator.

The findings in this report are not to be construed as an
official Department of the Army position unless so
designated by other authorized documents.

The contents of this report are not to be used for
advertising, publication, or promotional purposes.
Citation of trade names does not constitute an
official endorsement or approval of the use of such
commercial products.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM | | | | | | | | | |
|---|-----------------------|--|--|----------------|------------------|-----------|-----------------|---------|----------|-----------------|--------------------|
| 1. REPORT NUMBER | 2. GOVT ACCESSION NO. | 3. REPORT TYPE & CATALOG NUMBER | | | | | | | | | |
| Technical Report Y-84-1 | A142 590 | | | | | | | | | | |
| 4. TITLE (and Subtitle) | | 5. TYPE OF REPORT & PERIOD COVERED | | | | | | | | | |
| DELINEATION OF WETLAND BOUNDARIES USING VEGETATION WITHIN THE ALTAMAHA RIVER BASIN OF GEORGIA | | Final report | | | | | | | | | |
| 7. AUTHOR(s) | | 8. CONTRACT OR GRANT NUMBER(s) | | | | | | | | | |
| Boyd F. Vaughn, Jr., Robert J. Cooper, Joel H. Braswell, Robin Hart | | Contract No. DACW39-78-C-0092 | | | | | | | | | |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS | | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS | | | | | | | | | |
| Envirosphere Company 145 Technology Park Atlanta, Ga. 30092 | | Wetlands Research Program | | | | | | | | | |
| 11. CONTROLLING OFFICE NAME AND ADDRESS | | 12. REPORT DATE | | | | | | | | | |
| Office, Chief of Engineers, U. S. Army Washington, D. C. 20314 | | April 1984 | | | | | | | | | |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) | | 13. NUMBER OF PAGES | | | | | | | | | |
| U. S. Army Engineer Waterways Experiment Station Environmental Laboratory P. O. Box 631, Vicksburg, Miss. 39180 | | 70 | | | | | | | | | |
| | | 15. SECURITY CLASS. (of this report) | | | | | | | | | |
| | | Unclassified | | | | | | | | | |
| | | 15a. DECLASSIFICATION DOWNGRADING SCHEDULE | | | | | | | | | |
| 16. DISTRIBUTION STATEMENT (of this Report) | | | | | | | | | | | |
| Approved for public release; distribution unlimited. | | | | | | | | | | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | | | | | | | | | | |
| 18. SUPPLEMENTARY NOTES | | | | | | | | | | | |
| Available from National Technical Information Service, 5285 Port Royal Road, Springfield, Va. 22161. | | | | | | | | | | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) | | | | | | | | | | | |
| <table border="0"> <tr> <td>Altamaha River</td> <td>Sampling methods</td> </tr> <tr> <td>Community</td> <td>Transition zone</td> </tr> <tr> <td>Georgia</td> <td>Wetlands</td> </tr> <tr> <td>Plant community</td> <td>Wetland vegetation</td> </tr> </table> | | | | Altamaha River | Sampling methods | Community | Transition zone | Georgia | Wetlands | Plant community | Wetland vegetation |
| Altamaha River | Sampling methods | | | | | | | | | | |
| Community | Transition zone | | | | | | | | | | |
| Georgia | Wetlands | | | | | | | | | | |
| Plant community | Wetland vegetation | | | | | | | | | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) | | | | | | | | | | | |
| <p>This study develops a sound quantitative method for the assessment of the structure, location, and composition of transition zones adjacent to wetland communities in the Altamaha River Basin of Georgia. The resulting methods will be incorporated into a U. S. Army Corps of Engineers methodology for delineating wetlands.</p> | | | | | | | | | | | |

(Continued)

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. ABSTRACT (Continued).

~ Six sampling methods, three similarity indices, and four analytical strategies were investigated. Phase I of the study consisted of evaluation of sampling and analytical methods and selection of the best method. The best method proved to be the use of (a) 1- x 4-m contiguous quadrats for sampling along transects parallel to the moisture gradient and (b) Jaccard's community coefficients for data analysis.

Phase II of the study consisted of further evaluation of the best method. This method was found to be advantageous because:

- a. It does not rely on the use of indicator species; plant associations are used instead.
- b. It is uncomplicated, technically reproducible, and therefore amenable to the legal interpretation in defining wetland boundaries.
- c. The procedures are well documented in the literature.
- d. It is applicable to a variety of physiognomic types.
- e. Minimal training is required for proper application of the method by field personnel.

| | |
|---------------|--|
| Accession For | |
| NTIS GRA&I | <input checked="checked" type="checkbox"/> |
| DTIC TAB | <input type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification | |
| By _____ | |
| DTIC TAB | |
| ATC _____ | |
| Dist _____ | |
| A-1 | |



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

SUMMARY

of the Army, acting through the Corps of Engineers (COE), regulating the disposal of dredged and fill material into the United States, as a result of Federal Water Pollution Control Act of 1972 (PL 92-500). Critical to the implementation of any permit under Section 404 of PL 92-500 is the ability to locate the boundary and transition zone communities. The distinctive character of wetlands results from the interactions of vegetation and soils with hydrologic processes. All three of these factors should be considered when attempting to delineate boundaries. This study develops a sound quantitative method for the assessment of the structure, location, and composition of transition zones adjacent to wetland communities in the Altamaha River Basin of Georgia. Data obtained will be used in conjunction with soil and hydrologic techniques for wetland delineation.

Four different sampling methodologies, three similarity indices, and four strategies were investigated. All sampling methods involved random transects parallel to the wetness gradient.

Sampling and data analysis produced a "best method" for transitional determination using 1- by 4-m contiguous plots and Jaccard's coefficient to compare wetland and upland end quadrats to all other quadrats. All transects were analyzed individually to determine community boundaries. The unique and variable nature of the transition zones encountered was a significant factor affecting the choice of methodologies.

The "best method" described offers the following advantages:

1. It does not rely on the presence of "indicator" species; rather, the boundaries are ecologically determined by the position of the plant species comprising the wetland, upland, and transition zone communities.

2. It is uncomplicated, technically reproducible, and therefore amenable to the legal interpretation in defining wetland boundaries. The field and statistical procedures are well documented in the literature.

3. The "best method" is applicable to a variety of physiognomic types.

4. Field application is uncomplicated, and a minimum of training is required for proper implementation by field personnel.

PREFACE

At the request of the Office, Chief of Engineers, U. S. Army, the Environmental Laboratory (EL) of the U. S. Army Engineer Waterways Experiment Station (WES) conducted this research effort under the initial auspices of the Dredged Material Research Program (1978-1979) and later under the Dredging Operations Technical Support (DOTS) Program. This study is being published as part of the Wetlands Research Program (WRP). Technical monitors of the WRP for the Office, Chief of Engineers, were Dr. John R. Hall and Mr. Phillip C. Pierce.

The research was performed under Contract No. DACW39-78-C-0092 (entitled "Wetland Transition Zone Study Within the Altamaha River Basin of Georgia") by the Envirosphere Company of Atlanta, Ga. Authors of the report were Mr. Boyd F. Vaughan, Jr., Mr. Robert J. Cooper, Dr. Joel H. Braswell, and Dr. Robin Hart. Mr. Tom Mather provided valuable assistance in the field. Graphics were provided by Ms. S. L. Hull. Dr. Wilbur Duncan gave technical assistance in the identification of certain plant species. Dr. Robert Terry Huffman and Dr. Gary Tucker, Research Botanists, formerly of EL, developed the project's scope of work and provided valuable technical assistance throughout the study. Dr. D. R. Sanders, Sr., EL, was Leader, Wetlands Research Team, Wetland and Terrestrial Habitat Group, during the final review and publication of the report. The study was conducted under the general supervision of Dr. H. K. Smith, Environmental Resources Division (ERD), EL; Dr. C. J. Kirby, Chief, ERD, EL; and Dr. John Harrison, Chief, EL. Mr. Charles C. Calhoun was Program Manager, DOTS, and Dr. Smith was Program Manager, WRP.

The Commanders and Directors of WES during this study and the publication of this report were COL John L. Cannon, CE; COL Nelson P. Conover, CE; and COL Tilford C. Creel, CE. Technical Director was Mr. F. R. Brown.

This report should be cited as follows:

Vaughan, B. F., Jr., et al. 1984. "Delineation of Wetland Boundaries Using Vegetation Within the Altamaha River Basin of Georgia," Technical Report Y-84-1, prepared by Envirosphere Company, Atlanta, Ga., for the U. S. Army Engineer Waterways Experiment Station, CE, Vicksburg, Miss.

CONTENTS

| | <u>Page</u> |
|--|-------------|
| SUMMARY | 1 |
| PREFACE | 2 |
| CONVERSION FACTORS, U. S. CUSTOMARY TO METRIC (SI) UNITS OF MEASUREMENT | 4 |
| PART I: INTRODUCTION | 5 |
| Project Background | 5 |
| Description of Study Area | 6 |
| PART II: METHODS | 11 |
| Selection of Sampling Locations | 11 |
| Transect Location | 12 |
| Sampling Methodologies | 13 |
| Measures of Species Abundance | 15 |
| Data Analysis | 16 |
| Voucher Specimens | 18 |
| PART III: RESULTS AND DISCUSSION | 19 |
| Sampling Methodologies | 19 |
| Data Analysis | 24 |
| PART IV: CONCLUSIONS | 32 |
| REFERENCES | 34 |
| APPENDIX A: JACCARD COMMUNITY COEFFICIENTS FOR PHASE I SAMPLING . . . | A1 |
| APPENDIX B: JACCARD COMMUNITY COEFFICIENTS FOR PHASE II SAMPLING . . . | B1 |
| APPENDIX C: SPECIES DISTRIBUTION BY COMMUNITY FOR PHASE I SAMPLING | C1 |
| APPENDIX D: SPECIES DISTRIBUTION BY COMMUNITY FOR PHASE II SAMPLING | D1 |
| APPENDIX E: NATIONAL WETLANDS INVENTORY (NWI) EQUIVALENTS OF WETLAND TYPES USED IN THIS REPORT. | E1 |

CONVERSION FACTORS, U. S. CUSTOMARY TO METRIC (SI)
UNITS OF MEASUREMENT

U. S. customary units of measurement used in this report can be converted to metric (SI) units as follows:

| <u>Multiply</u> | <u>By</u> | <u>To Obtain</u> |
|-----------------------|------------|-----------------------------|
| acres | 0.4046873 | hectares |
| cubic feet per second | 0.02831685 | cubic metres per second |
| Fahrenheit degrees | 5/9 | Celsius degrees or Kelvins* |
| feet | 0.3048 | metres |
| inches | 2.54 | centimetres |
| miles (U. S. statute) | 1.609347 | kilometres |
| square miles | 2.589998 | square kilometres |

* To obtain Celsius (C) temperature readings from Fahrenheit (F) readings, use the following formula: $C = (5/9)(F - 32)$. To obtain Kelvin (K) readings, use $K = (5/9)(F - 32) + 273.15$.

WETLAND BOUNDARIES USING VEGETATION WITHIN THE ATLANTA RIVER BASIN OF GEORGIA

PART I: INTRODUCTION

Project Background

Department of the Army, acting through the Corps of Engineers
able for regulating the disposal of dredged and fill material
United States, as a result of the Federal Water Pollution
Acts of 1972 (PL 92-500). Until 1975, however, the regula-
tion was directed toward only navigable waters. Since that time, new
law has provided for extending jurisdiction into all waters of
the United States, including wetlands. Such jurisdiction is essential to
maintain the chemical, physical, and biological integrity of the Nation's

to the implementation of any permit program under Sec-
tion 404 of the Clean Water Act. Current COE rules and reg-
ulations (Register, 19 July 1977) define wetlands as:

Wetlands are areas that are inundated or saturated by surface or
groundwater at a frequency and duration sufficient to support,
under normal circumstances do support, a prevalence
of plants typically adapted for life in saturated soil
conditions. Wetlands generally include swamps, marshes, bogs,
and similar areas.

Wetland determination, determination of an area as wetland is contingent
upon the relationship of soils and vegetation with the hydrologic regime.
Wetlands are bordered by an area of transition from wetland to up-
land. A problem exists in that the boundary of this transition
is not clearly defined, since it shares species common to both sides of
the boundary. Determination of a wetland boundary requires examina-
tion of the transition zone and its relation to the wetland areas. It was the
purpose of this study to formulate a methodology for delineating the bound-
ary, defining the transition zones present, and distinguishing
wetlands from uplands.

The study develops a sound quantitative methodology for the assess-
ment of the location and composition of transition zones adjacent to wetland

plant communities in the Altamaha River Basin of Georgia in an area extending from the Atlantic coast inland to Doctortown, Georgia. Wetlands boundaries can be defined by the change in species composition along the moisture gradient. This study compared different sampling methods to determine the most accurate and cost-effective procedure for delineating wetlands. Wetland types differing in physiognomy, salinity, degree of disturbance, and transition zone width were investigated to evaluate the general applicability of the different methodologies. The physiognomic types studied were a tree-dominated wetland, shrub-dominated wetland, graminoid marsh, flat, and open-water habitat with macrophytic vegetation.

5. The field sampling effort consisted of two phases. Phase I was the initial field application of the different methodologies, while Phase II was a field verification of the "best" methodology.

Description of Study Area

Location

6. The study area is composed of the Altamaha River Basin from the Atlantic coast inland to Doctortown, Georgia, including St. Simons Island and the system of estuaries and marshlands between the island and the coast. The Altamaha River, the headwaters of which originate in the Piedmont, is one of the two major river systems of the Atlantic drainage in Georgia. The Altamaha has the greatest discharge of any river along the Georgia coast. It has a drainage area of approximately 13,600 square miles* and an average discharge of 12,600 cfs (U. S. Geological Survey 1959). Several miles from the coast it divides into four distributaries (Darien, Butler, Champney, and South Altamaha Rivers) which empty into Altamaha and Doboy Sounds.

7. From the coast inland, the Altamaha is bordered by abandoned rice fields (to mile 13) which have a general elevation of 7 ft above low water. The general slope of the Altamaha is 0.7 ft per mile, but there are short segments less than 1 mile long where slopes reach 3 and 4 ft per mile. The sharpest curve has a radius of 200 ft and there are 50 curves with radii less than 500 ft (Wharton 1978).

Physiography and geology

8. Soils of the Piedmont Province are derived from pre-Cambrian

* A table of factors for converting U. S. customary units of measurement to metric (SI) units is provided on page 4.

crystalline rocks. Marking the border between the Piedmont and Coastal Plain is the Fall Line, indicating the farthest advance of the sea late in the Mesozoic Era. The Coastal Plain is overlaid by many deposits caused by changes in sea level in the Tertiary and Quaternary Periods, with the most recent deposits located at the coast.

9. Formation of the barrier islands has been explained by several theories, the most commonly accepted of which attributes the phenomenon to dune formation, followed by severe flooding, which occurred at the end of the Pleistocene ice ages. The area landward of the largest and most stable dunes would therefore become a lagoon, eventually filling with sediment and evolving into a salt marsh. The coastal islands are of two ages. The older islands (e.g., St. Simons) were formed during the Silver Bluff Pleistocene submergence (25,000-36,000 years ago) described above. The newer islands (e.g., Sea Island) were formed 4000-5000 years ago during the Holocene (Hoyt 1967, 1968).

10. The lower Georgia Coastal Plain was developed as a result of a series of barrier island formations. Each series was followed by a recession of sea level, resulting in a series of terraces. The Wicomico shoreline, the oldest such terrace, was as much as 100 ft higher than the present shoreline. Subsequent barrier island formations were the Penholoway (75 ft above present sea level), Talbot (40-45 ft), Pamlico (25 ft), Princess Anne (15 ft), Silver Bluff (5 ft), and Holocene (Hoyt and Hails 1967).

11. Tidal marshes are formed in conjunction with barrier island development. The areas landward of the partially submerged dunes are subject to less disturbance than open waters, permitting clay and silt sediments to be deposited. This permitted extensive communities of salt-tolerant marsh plants, such as smooth cordgrass (*Spartina alterniflora*) to develop. Tidal action continually contributes new sediments at a very slow rate, and receding tides form an extensive drainage system of tidal creeks and rivers, creating a network of marsh islands throughout the estuarine system.

Climate

12. The coastal region of Georgia has a relatively moderate climate, with slightly lower average temperatures than on the mainland due to offshore winds. Winters are relatively short and mild. Mean minimum and maximum January temperatures are 44° F and 64° F, respectively. Mean minimum and maximum July temperatures are 72° F and 90° F, with mean annual rainfall approximately 52 in. Rain is most frequent in summer and early fall. The driest period is

November through February (Carter 1967). Tropical storms are common between August and October, yet in the period 1886-1968 only 56 reached hurricane magnitude and few have caused serious damage (Carter 1970).

Soils

13. Wetland soils are very poorly drained and are flooded periodically if not regularly. Largely a result of alluvial deposits, wetland soils are predominantly clays or silty clays, although they also contain some fine sand.

14. Swamp soils, associated with freshwater swamp forests, are located at higher elevations and are seldom covered by tidal waters. They are generally flooded in winter and spring and receive fresh deposits of sediments. The surface layer is generally black loam or clay loam, overlaying mostly dark-gray clay (Byrd et al. 1961).

15. Tidal marsh soils are covered twice daily by tides. Johnson et al. (1974) have summarized the sources of the sediments in the salt marsh as derived either from the continental shelf, the mainland rivers, the marsh itself, or organic deposits. Tidal marshes depend on upland freshwater sources for much of their nutrient and biological richness (Gosselink, Odum, and Pope 1973). Haines (1975) felt that substantial amounts of trace minerals, silicates, and organic nutrients could be contributed by freshwater discharge into tidal areas. The upper few centimetres of salt-marsh soils are usually brown. The lower portion is usually black with a nearly neutral pH. Sulfides, methane, and ferrous compounds are present due to the anaerobic conditions (Wharton 1978). High marsh sites tend to have more sand and less clay and organic matter than low marsh sites (Teal and Kanwisher 1961).

16. Wet alluvial soils (Byrd et al. 1961) occur at the mouth of the Altamaha River. They are covered by salt water at high tides and are swept by fresh water when the river is high. These soils are similar to tidal marsh soils but include periodic lenses of sand. The distribution of grain size is such that the coarser materials are located in the estuary and offshore delta areas, and finer materials accumulate in the tidal flats (Visser and Howard 1974).

17. The soils of St. Simons Island are characteristically strongly acidic, low in fertility, and either excessively drained (Lakewood Series) or poorly drained (Leon, Rutledge, Plummer Series).

Flooding history

18. Wetland plant communities are largely determined by the duration,

periodicity, and depth of flooding. On the Altamaha, flooding originates from two sources, river floodstage and tidal action.

19. The U. S. Soil Conservation Service has estimated that tidal marsh acreage in Glynn and McIntosh Counties is 78,936 and 83,132, respectively. Tidal waters inundate marsh areas twice daily, with tidal influence extending far upriver. The limit of inland tidal influence on the Altamaha River is estimated to be river mile 42 near the Long-McIntosh County line. The limit of salinity influence is located at approximately river mile 18.9 (Georgia Department of Natural Resources 1976). Thus some areas upriver experience daily water level fluctuations of fresh or brackish water, while others are affected only when spring or storm tides, augmented by onshore winds, inundate the floodplain with up to several feet of water. Similarly, higher areas of salt marsh, characterized by needlerush (*Juncus roemerianus*), are inundated with salt water only during extremely high tides.

20. Spring floods on the Altamaha inundate and flush the floodplain, providing the area with new sediments as well as transporting sediments to the estuary and beyond. River swamps are generally flooded by 1 or more feet of water for 6 months or more of the year (Bozeman and Darrell 1975). By holding for a time these overbank river flows, swamps may also increase shallow groundwater supplies. In times of dry weather, the shallow watertable aquifer feeds the river swamp (Georgia Department of Natural Resources 1976).

Vegetation

21. Wetland vegetation of the study area may be categorized as either swamp or marsh. Swamps are typically hardwood swamps associated with fresh water. Marshes may be either salt or fresh and are typically covered with herbaceous vegetation.

22. River swamp ecosystems are highly diverse, with the dominant species being water tupelo (*Nyssa aquatica*) and bald cypress (*Taxodium distichum*) in extremely hydric conditions and overcup oak (*Quercus lyrata*), green ash (*Fraxinus pennsylvanica*), Carolina ash (*Fraxinus caroliniana*), and black gum (*Nyssa biflora*) in slightly less hydric conditions (Bozeman and Darrell 1975). Klawitter (1962) and Applequist (1959) also include red maple (*Acer rubrum*) as a major component of these communities.

23. Freshwater marshes occur primarily near the mouth of the Altamaha and extend for some distance upriver before being replaced by swamp forest. Much of the freshwater marsh area was once swamp forest that was cleared and

diked for the growing of rice. Shallow freshwater marshes contain a variety of species. The deeper freshwater marshes are more extensive, occupying an area of approximately 25,000 acres along the Georgia coast (Johnson et al. 1974). Sawgrass (*Cladium jamaicense*) is a dominant species in much of this area. Cattails (*Typha* spp.) are common along the deeper margins. As salinities increase to brackish conditions, giant cutgrass is replaced by big cordgrass (*Spartina cynosuroides*) and salt-marsh bulrush (*Scirpus robustus*).

24. Zonation in the salt marsh is primarily related to elevation, as it determines frequency, depth and duration of inundation, and soil salinity (Teal and Teal 1969). The most extensive monospecific marshes consist of smooth cordgrass, covering an area of 285,650 acres (Spinner 1969). Higher, infrequently flooded salt marshes contain needlerush.

PART II: METHODS

Selection of Sampling Locations

25. The study area consisted of the Altamaha River Basin inland to Doctortown, Georgia. Access problems and time considerations dictated that all sampling sites be located in the lower half of this area. The results are applicable to the entire basin since no major wetland types or salinity conditions were excluded by this approach. Potential sites were selected on the basis of access, vegetation homogeneity, and how well they represented the desired physiognomic type. Color-infrared aerial photographs and topographic maps (scale 1:24,000) were used to identify potential sites. Appropriate State agencies were consulted and a reconnaissance trip was made to each site before sampling commenced. A COE representative visited all sites prior to Phase I and II sampling and was given an opportunity to make recommendations concerning the candidate areas. Sampling locations are presented in Figures 1 and 2.




Figure 1. Sampling location for Glynn County, Georgia




Figure 2. Sampling locations for McIntosh County, Georgia

Transect Location

26. Sampling area boundaries were established based on homogeneity of vegetation and accessibility. Transects were randomly located. Within a suitable sampling area there was an infinite number of transect locations. By breaking the shoreline into a finite number of nonoverlapping, 4-m-wide transects, the transects could be numbered and then selected randomly. Five transects were selected in this manner and were placed parallel to the moisture gradient with the aid of a compass. Gradient analysis requires that sampling units be parallel to the gradient being analyzed.

27. The first quadrat of each transect was placed at approximately the same distance into the wetland community. (In this report, quadrat refers to either a square or rectangular sampling area.) This was accomplished using a common point of reference for all transects. The point was defined as the uplandmost edge of the dominant wetland vegetation present. The transect followed the appropriate compass heading until a zone was reached which supported vegetation known not to tolerate saturated or flooded conditions for an extended period of time. Transect length ranged from 8 to 25 m, depending primarily on transition zone width. The same procedure was followed for both Phase I and II sampling.

Sampling Methodologies

28. All sampling was conducted by a project team consisting of two biologists. Five different sampling methodologies were examined during the Phase I sampling: (a) square quadrats randomly spaced along the transect, (b) square quadrats systematically spaced along the transect, (c) rectangular quadrats randomly spaced along the transect, (d) rectangular quadrats systematically spaced along the transect, and (e) line intercept. The quadrat spacing was dictated to a certain degree by the length of the transect being sampled. An attempt was made to place at least two quadrats in each vegetation zone (i.e., wetland, transition, and upland). However, the transition zones' narrow width often restricted the number of quadrats that could be placed within it. Quadrat size chosen is that customary for the stratum being considered and is presented in Table 1 (Mueller-Dombois and Ellenberg 1974).

Table 1
Sample Unit Sizes Used During Phase I and Phase II

| Communities | Strata | Sample Unit Size, m | | |
|------------------------------|------------|---------------------|---------|-----------|
| | | Line Intercept | Square | Rectangle |
| Phase I | | | | |
| Salt Flat | Herbaceous | 4 | 1 × 1 | 1/2 × 2 |
| Disturbed Fresh water | Shrub | 4 | 4 × 4 | 2 × 8 |
| Marsh, Steep | Canopy | 4 | 10 × 10 | 5 × 20 |
| Swamp Forest (two transects) | | | | |
| Gradual Swamp | Herbaceous | 1 | 1 × 1 | |
| Forest, Shrub | Shrub | 1 | | 1 × 4 |
| Open Water, Freshwater | Canopy | 1 | | 1 × 4 |
| Marsh, Saltwater | | | | |
| Marsh, Steep Swamp | | | | |
| Forest | | | | |
| Phase II | | | | |
| All Communities | Herbaceous | | 1 × 1 | |
| | Shrub | Not Used | | 1 × 4 |
| | Canopy | | | 1 × 4 |

29. Transition zone width also imposed constraints on quadrat size. Sampling areas had to be large enough to include a representative sample of the most abundant species within the zone, yet be small enough so that each quadrat was fairly homogeneous and changes along the gradient were not obscured. Moreover, quadrat sizes used in transition zones set the size for quadrats in upland and wetland zones since valid comparisons of successive samples cannot be made between quadrats of unequal size. These five methods with the described sampling unit sizes were used in the first three communities sampled (salt flat, disturbed freshwater marsh, and two transects in the steep swamp forest). The tree canopy in the salt flat upland was measured in shrub plots because the upland was an island with insufficient area for replicate 100-m² plots. The line intercept method involved use of 4-m segments as the basic sampling unit during Phase I. This sampling methodology is described by Canfield (1941).

30. After sampling three communities, adjustments were made in the sampling methodology. The shrub, open water, freshwater marsh (undisturbed), saltwater marsh, gradual swamp forest, and steep swamp forest communities were sampled using both contiguous 1- by 4-m quadrats (1-m side parallel to the moisture gradient) with nested 1- by 1-m quadrats and the line intercept method. The 1- by 4-m quadrat size represents a compromise between obtaining the resolution desired in locating the transition zone boundary and assuring an adequate vegetation sample.

31. A transect was established by extending two 100-m tapes 4 m apart parallel to the moisture gradient from wetland to upland. Additional tapes were extended between the 100-m tapes at 1-m intervals to establish a 1- by 4-m sample plot. Trees and shrubs were sampled in the resulting sample plot, and herbaceous species were sampled in two 1- by 1-m plots adjacent to the longer tapes. This procedure was repeated for each 1-m interval along the transect.

32. The same 100-m tapes identified above also served as the base lines for sampling units used for the line intercept method. All species intercepting the tapes with a leaf or stem or vertical projection of a leaf or stem were included in the sampling. The time required to sample by each method was recorded, and the order of sampling was alternated in consecutive transects so that familiarity with transect vegetation would not bias the time each method required.

33. Elevations for Phase I were measured with a hand level and telescoping level rod. Measurements were taken every 1 m on each of the five transects sampled per wetland type. Elevations were measured to the nearest 0.05 ft. Where possible, elevations along a particular transect were taken from a single location. All elevations were relative, with the first elevation in the wetland representing zero. Elevations during Phase II sampling were taken only in the first and last quadrat, giving the total change in elevation for a transect.

Measures of Species Abundance

34. The abundance measure should be a sensitive indicator of the "importance" of a species. This measure helps delineate a zone or zones along a gradient where the species most successfully competes in comparison with other species present in the community.

35. Three measures of species abundance were considered during Phase I of the sampling effort: percent cover, diameter at breast height (canopy strata only), and stem counts (shrub strata only). Percent cover is defined as the percent of the ground within a quadrat intercepted by a projection of the canopy. Cover values were recorded for all herbaceous and shrub species rooted in the plot with an estimated cover of 5 percent or greater. All values were estimated to the nearest 5 percent. Diameters at breast height (dbh) were taken for the canopy stratum, while stem counts were made in the shrub stratum. This approach assured a certain flexibility during the data analysis phase at which time techniques using scaled values or a presence-absence representation could be investigated, since these required no additional field sampling.

36. Project team members made independent estimates of cover. Estimates usually agreed within 5 percent, with maximum differences seldom exceeding 10 percent. The strata measured were defined as follows:

- a. Herbaceous stratum. All herbaceous and woody species equal to or less than 1 m in height.
- b. Shrub stratum. Woody species with heights greater than 1 m and diameters (dbh) less than 5 cm.
- c. Canopy stratum. Woody species with diameters (dbh) greater than or equal to 5 cm.

37. Phase II sampling used only percent cover as the measure of species abundance. Cover value estimates were made for all species with an estimated

cover of 5 percent or greater. Species which were not rooted in but had cover in a plot were included if the projection of the quadrat perpendicular to the moisture gradient would have included the plant.

Data Analysis

38. Beta diversity (BD) is defined as between-habitat diversity, or the degree of change in species composition of communities along a gradient (Whittaker 1975). Three different measures of beta diversity were investigated during the analysis of Phase I data.

39. Jaccard's (1901, 1912, 1928) community coefficient (CC) is based on the presence-absence relationship between the number of species common to two areas and the total number of species present:

$$\text{Jaccard CC} = \left(\frac{c}{a + b + c} \right) 100$$

where: c = the number of species common to both areas
 a = the number of species unique to area a
 b = the number of species unique to area b

40. A similar approach but one which incorporates importance values (percent cover for this report) for the various species is presented by Bray and Curtis (1957):

$$\text{Percent Similarity (PS)} = \sum_{i=1}^n \text{Min} (P_{ij})(P_{ik})$$

where P_{ij} and P_{ik} are the relative abundance values for species i in quadrats j and k (the quadrats being compared), and n is the number of species common to both quadrats. Relative abundance values were calculated in the following manner:

$$\text{Relative abundance for species i} = \frac{\text{Percent cover for species i in quadrat j}}{\text{Total percent cover for all species in quadrat j}}$$

This method gives more weight to the dominant species where the Jaccard community coefficient gives equal weight to all species.

41. Bratton (1975) offers this dynamic definition of BD:

$$B(g) = \sum_{i=0}^n B_i(g)$$

Where $B(g)$ is BD at a point along the gradient, and $B_i(g)$ is the rate at which abundance of the i^{th} species is changing at point g . The total number of species present at point g is represented by n . This measure was approximated by summing the absolute change in abundance (i.e., cover values) of each species between two successive quadrats. The relative rate of change was calculated by dividing $B(g)$ by the total species cover in the first of the two quadrats. A significant increase in the relative rate of change along the gradient should represent a change in community structure.

42. Two alternative strategies were utilized to delineate transition zone boundaries using the three BD indices described above. The first strategy involved calculating BD indices for successive pairs of quadrats. Thus 1 was compared to 2, 2 to 3, 3 to 4, etc. All three indices were examined using this approach. A second alternative compared all quadrats to the two end quadrats. Indices were calculated comparing 1 to 2, 1 to 3, etc., until 1 had been compared to all quadrats along the transect. Given a transect with 20 quadrats, the procedure was then reversed comparing 20 to 19, 20 to 18, and finally 20 to 1. Jaccard's CC along with percent similarity (Bray and Curtis 1957) were calculated using this second alternative.

43. Communities were analyzed both by combining the five transects into a composite transect and by analyzing individual transects. Species cover values for the composite were arrived at by averaging all cover values for a given species in a given quadrat for all five transects.

44. The different combinations of analytical strategies investigated are presented in Table 2. In certain instances analytical techniques were tried on only a portion of the data.

45. Mean, variance, standard deviation, and standard error of the mean were calculated for each species' cover values for a particular quadrat over the five transects or ten line intercepts sampled. Similarly numbered quadrats within a particular community were also compared using percent similarity. These simple statistics gave an indication of the between-transect variability in each community.

Table 2
Alternative Analytical Methods Investigated

| | Jaccards' Community Coefficient | Percent Similarity (Bray and Curtis 1957) | Beta Diversity (Bratton 1975) |
|--------------------------|---------------------------------------|---|----------------------------------|
| <u>Phase I</u> | | | |
| Successive Quadrats | * | * | * |
| Quadrats to End Quadrats | * | * | |
| Composite Transect | ** | * | * |
| Individual Transect | * | ** | |
| <u>Phase II</u> | | | |
| Successive Quadrats | * | * | |
| Quadrats to End Quadrats | * | * | |
| Composite Transect | | * | |
| Individual Transect | * | | |

* All communities were analyzed.

** Selected communities were analyzed.

46. The analysis of Phase II data concentrated on a further investigation of Jaccard's community coefficient.

Voucher Specimens

47. Voucher specimens were collected of all dominant plant species in all communities sampled. The plants were pressed, dried, and sprinkled with paradichlorobenzene to prevent insect damage. Each plant was labeled as to scientific name, collection location, collector, and date of collection. Nomenclature followed Radford, Ahles, and Bell (1968) with updating by Wilbur H. Duncan, plant taxonomist at the University of Georgia.

PART III: RESULTS AND DISCUSSION

Sampling Methodologies

Quadrat shape and size

48. Data were analyzed after sampling three communities (salt flat, disturbed freshwater marsh, steep swamp forest). Significant differences between data from square and rectangular plots occurred only where a species was restricted to a narrow zone of the moisture gradient relative to plot size. *Nyssa biflora*, for example, was restricted to a narrow zone at the juncture of the open marsh with a wooded floodplain. This species was virtually absent in the rectangle, but did occur in the upper half of the square in the wetland zone (Table 3). *Liquidambar styraciflua* and *Pinus taeda* were more abundant in the upper half of the transition and upland zones, respectively, and were therefore both better represented in the square plots. The rectangular quadrats more closely defined vegetation zone boundaries where zones were narrow because they bordered shorter segments of the moisture gradient. However, as Tables 3, 4, and 5 illustrate, there were no significant differences between square and rectangular quadrats in number of woody individuals included or mean percent cover of herbaceous species.

49. The line intercept method did not differ greatly from the square and rectangular quadrats in the percent cover recorded for herbaceous species (Table 4). However, the number of woody plants included in the line intercept method was significantly (at least a factor of two and often much greater) lower than in either quadrat shape and did not adequately represent these species (Table 5). A much larger number of line intercepts would be required to get data equivalent to that obtained in a 10- by 10-m plot.

50. Time required for sampling square and rectangular quadrats did not differ greatly (Table 6). The line intercept method required approximately the same time in the predominantly herbaceous salt flat but less time in the disturbed freshwater marsh which had wooded transition and upland zones. This difference in sampling time can be attributed to the different number of trees present in the two communities. A large number of trees increased sampling time for rectangles and squares, while not significantly increasing the time required for the line intercept method due to the smaller number of trees included in the latter survey.

Table 3

Abundance of Tree Species in Square and Rectangular Quadrats in the Disturbed Freshwater Marsh

| Species | Zone* | Square Quadrats | | Basal Area, cm ² Mean Values | Rectangular Quadrats | | Basal Area, cm ² Mean Values |
|--|-------|-----------------|-------|--|-------------------------|-------|--|
| | | Frequency | Stems | | Frequency | Stems | |
| <i>Nyssa biflora</i> | WL** | 4 | 3.2 | 348 | 1 | 0.8 | 31 |
| | TZ | 5 | 3.6 | 958 | 5 | 5.2 | 1029 |
| <i>Myrica cerifera</i> | TZ | 4 | 1.2 | 30 | 4 | 1.0 | 35 |
| <i>Liquidambar styraciflua</i> | TZ** | 3 | 1.2 | 301 | 3 | 0.6 | 88 |
| | UL | 4 | 1.0 | 127 | 3 | 1.0 | 184 |
| <i>Pinus taeda</i> | UL** | 4 | 2.0 | 1058 | 3 | 1.4 | 490 |
| <i>Quercus hemispherica</i> | UL | 3 | 1.2 | 637 | 4 | 2.8 | 856 |
| <i>Quercus nigra</i> - <i>hemispherica</i> (hybrid) | UL | 4 | 2.2 | 1044 | 4 | 2.4 | 1644 |

* WL = Wetland, TZ = Transition Zone, UL = Upland.

** Significant difference ($\alpha = 0.05$) between square and rectangular quadrats in at least one measure.

Table 4

Mean Percent Cover (\pm SEM) of Four Herbaceous Species and One Shrub Species

| Community | Species | Zone* | Square | | Rectangle | | Line Intercept |
|----------------------------|---------------------------------|-------|-----------------|-----------------|-----------------|-------|----------------|
| | | | Frequency | Stems | Frequency | Stems | |
| Disturbed Freshwater Marsh | <i>Cladium jamaicense</i> | WL | 45.0 \pm 11.0 | 35.0 \pm 14.0 | 36.0 \pm 17.0 | | |
| Disturbed Freshwater Marsh | <i>Pontederia cordata</i> | WL | 7.0 \pm 1.0 | 5.0 \pm 0.0 | 8.0 \pm 1.0 | | |
| Disturbed Freshwater Marsh | <i>Eleocharis quadrangulata</i> | WL | 33.0 \pm 10.3 | 37.0 \pm 9.8 | 25.0 \pm 8.0 | | |
| Salt Flat | <i>Salicornia virginia</i> | WL | 21.0 \pm 7.0 | 19.0 \pm 3.0 | 15.0 \pm 2.0 | | |
| Steep Swamp Forest | <i>Myrica cerifera</i> | WL | 7.0 \pm 1.0 | 10.0 \pm 3.0 | 1.0 \pm 1.0 | | |

* WL = Wetland.

Table 5
Woody Individuals Included in Squares, Rectangles, and
Line Intercept Sampling Units

| Community | Zone* | Square | Rectangle | Line Intercept |
|-----------------------------------|-------|------------------|----------------|-----------------|
| | | Mean \pm SEM** | Mean \pm SEM | Mean \pm SEM |
| <u>Disturbed Freshwater Marsh</u> | WL† | 4.4 \pm 1.9 | 0.8 \pm 0.8 | 0.6 \pm 0.4 |
| | TZ | 8.6 \pm 1.4 | 9.3 \pm 1.3 | 0.4 \pm 0.2 |
| | UL | 9.4 \pm 1.0 | 8.4 \pm 1.2 | 0 |
| <u>Steep Swamp Forest</u> | WL | 16.0 \pm 0.0 | 18.5 \pm 3.5 | 2.3 \pm 0.3 |
| | TZ | 17.0 \pm 6.0 | 18.5 \pm 4.5 | 3.5 \pm 0.7 |
| | UL | 21.0 \pm 4.5 | 12.5 \pm 2.5 | 5.3 \pm 0.75 |
| <u>Salt Flat</u> | UL | 0 | 0 | 0 |
| | TZ | 12.7 \pm 2.4 | 15.0 \pm 5.3 | 0.25 \pm 0.25 |
| | UL | 13.8 \pm 0.8 | 17.5 \pm 2.5 | 0 |

* WL = Wetland, TZ = Transition, UL = Upland.

** SEM = standard error of the mean.

† Significant difference ($\alpha = 0.05$) between rectangle and square.

Table 6
Time Required to Sample All Strata

| Community | Sampling Method (minutes \pm SEM) | | |
|-----------------------------------|-------------------------------------|------------|----------------|
| | Square | Rectangle | Line Intercept |
| <u>Disturbed Freshwater Marsh</u> | 12 \pm 4 | 16 \pm 2 | 5 \pm 1 |
| <u>Salt Flat</u> | 11 \pm 2 | 7 \pm 2 | 10 \pm 6 |

Preliminary results indicated that adjustments should be made in the sampling methodology. Random and systematic placement of quadrats were both inappropriate because of the narrow transition zones present and were eliminated from future sampling. Surveys taken in a 400-m² plot in the gradual swamp forest, a 250-m² plot in the freshwater marsh, and 100-m² plots in the disturbed freshwater marsh and steep swamp forest showed that the average tree density in the wooded areas of these communities was one individual per 10 m². However, the plot size required to adequately sample trees was too large to obtain the resolution desired in locating the transition zone boundaries. Therefore it was decided that the original quadrat sizes, while appropriate for sampling the various strata, would result in an unacceptable loss of resolution.

51. The five remaining communities plus the steep swamp forest were

sampled using both contiguous 1- by 4-m quadrats and the line intercept method. Preliminary results had indicated that the line intercept method did not adequately sample the canopy but still might be appropriate for measuring the herbaceous and shrub strata. The line intercept (sample unit size) was reduced from 4 to 1 m.

52. The most favorable sampling method was that which, considering the time spent sampling, gave the highest frequency of occurrence of species among the replicate transects and showed the lowest variability using the ratio of the standard error of the mean (SEM) to the mean as a variability indicator. The frequency of occurrence of woody species was highest in the 1- by 4-m quadrats (35). Herbaceous species were more frequently observed in the 1- by 1-m quadrats (25), while the line intercept method (5) was lower by a factor of five. The frequency data are presented by community in Table 7.

Table 7
Frequency of Occurrence of Woody and Herbaceous Species

| Community | Method | | | | |
|----------------------|----------------|-------|------------------|-------|------------------|
| | Line Intercept | | 1- × 1-m quadrat | | 1- × 4-m quadrat |
| | Shrubs | Herbs | Shrubs | Herbs | |
| Freshwater Marsh | 1 | 1 | 1 | 12 | 4 |
| Gradual Swamp Forest | 1 | - | 1 | 0 | 6 |
| Steep Swamp Forest | 1 | 1 | 0 | 0 | 9 |
| Open Water | 1 | 2 | 1 | 6 | 4 |
| Saltwater Marsh | 0 | 1 | 1 | 4 | 6 |
| Shrub | 0 | 0 | 0 | 3 | 6 |
| Total | 4 | 5 | 4 | 25 | 35 |

53. The ratio of the SEM/mean for species which had a frequency of two or more was determined for each sampling method. A low ratio indicates low variability. Twenty shrub species had the lowest ratio in 1- by 4-m quadrats, while one species was lowest in line intercept segments (Table 8). Twelve herbaceous species showed the lowest ratio in 1- by 1-m quadrats, and two were lowest in the line intercept segments. Time differences observed during preliminary sampling were not observed during this latter stage of Phase I (Table 9) since fewer trees were included in the 1- by 4-m quadrats than in the 10- by

Table 8
Number of Woody and Herbaceous Species Showing the Lowest Variability (SEM)

| Community | Method | | | | |
|----------------------|----------------|-------|------------------|-------|-------------------|
| | Line Intercept | | 1- × 1-m quadrat | | 1- × 4-m |
| | Shrubs | Herbs | Shrubs | Herbs | quadrat Shrubs |
| Freshwater Marsh | 0 | 2 | 0 | 7 | 5 |
| Gradual Swamp Forest | 1 | - | 0 | - | 2 |
| Open Water | 0 | 0 | 0 | 3 | 4 |
| Shrub | 0 | 0 | 1 | 2 | 5 |
| Steep Swamp Forest | 0 | - | 0 | - | 5 |
| Saltwater Marsh | 0 | - | 1 | - | 2 |
| Total | 1 | 2 | 2 | 12 | 20 |

Table 9
Sampling Time Required, in Minutes

| Community | Quadrat (Mean + SEM) | Line Intercept (Mean + SEM) |
|----------------------|-------------------------|--------------------------------|
| Freshwater Marsh | 34±3 | 34±1 |
| Gradual Swamp Forest | 20±2 | 18±1 |
| Open Water | 14±2 | 15±0 |
| Shrub | 15±2 | 16±2 |
| Saltwater Marsh | 22±2 | 29±2 |
| Steep Swamp Forest | 18±1 | 22±2 |

10-m quadrats, thus reducing the time spent measuring trunk diameters.

54. Since the 1- by 4-m quadrats sampled more woody species with a lower SEM/mean ratio than the line intercept method in approximately the same time, this method was judged superior for sampling the shrub and tree strata. The 1- by 1-m quadrat was the better sample size for herbaceous species for the same reasons. Therefore contiguous 1- by 4-m plots with nested 1- by 1-m quadrats was the "best" sampling method and was used in Phase II sampling.

Abundance measures

55. Separate analysis of strata resulted in the loss of valuable

information since transition zones proved to be areas of physiognomic as well as floristic change. A shift from predominantly herbaceous species to shrub and tree species comprised a critical part of some transition zones which could be characterized only if all strata were combined in the analysis. Two options were considered. First, an abundance measure could be employed that would represent all physiognomic forms. Or second, different abundance measures could be used for each stratum and standardized to an index of relative abundance before similarity comparisons were made.

56. Percent cover was the most flexible of the abundance measurements considered. It can be determined rapidly and represents the ability of a species to exploit space. Since species of all strata share this biological attribute, percent cover is appropriate for measuring all physiognomic forms. By using percent cover, the investigators retained the flexibility of using cover estimates or a simple presence/absence representation during data analysis.

57. Therefore, it was decided that percent cover was the most appropriate abundance measure for use in this study. Cover values for all strata were considered equal in "importance." A species' total cover in a quadrat was recorded as one measurement, even though it might appear in several different strata.

Data Analysis

58. The manner in which community composition changes along an environmental gradient and the manner in which different communities intergrade have resulted in several opposing schools of thought on the subject. For example, the "organismal" school holds that there is discontinuous variation in community composition along environmental gradients and that communities exhibit sharp boundaries along such gradients. The individualistic school holds that community composition varies gradually along environmental gradients, with continuous intergradation between communities (Cox 1980). Whittaker (1975), in his discussion of plant responses to elevational gradients, defines the principle of species individuality:

Each species is distributed in its own way, according to its own genetic, physiological, and life cycle characteristics and its way of relating to both physical environment and interactions with other species; hence no two species are alike in distribution.

Similar responses were observed in this study. The observations of this study further agree with the principle of community continuity:

The broad overlap and scattered centers of species populations along a gradient imply that most communities intergrade continuously along environmental gradients, rather than forming distinct, clearly separated zones.

59. In his work in the Great Smoky Mountains, Whittaker concluded that competition did not usually produce sharp boundaries between species populations and that evolution of species in relation to one another does not produce well-defined groups of species with similar distributions. Although discrete boundaries were sometimes observed by him and by the authors of this study, the causes of such boundaries are frequently explained by a sharp environmental discontinuity or by human disturbance. Also, pure stands of one species (e.g., *Juncus roemerianus*) may form a distinct community boundary. Overall, the authors of this study view the environmental gradient from wetland to upland as a line of continuously changing species that seldom exhibits a discrete, easily recognized boundary between associations.

60. Therefore in any classification system, the boundaries between community types will be more or less arbitrary, for these boundaries are determined by the characteristics chosen for classification and the ecologist's choice of where to place the boundaries. There is no single correct way to classify communities. A number of different classification systems have developed. Species dominance, strata structure, and species composition are among the more common community characteristics employed, with species composition being the most common. Species composition in the form of either percent cover or a presence-absence representation was the principal characteristic used for boundary determination in this study. The methods presented in Table 2 were those used for data analysis.

61. Not all of the species sampled were used in the analysis. Species present in both the first and last quadrat were excluded since these plants were not responding to the moisture gradient in an observable way and thus could not contribute to determining the transition zone boundary. Species appearing only once were also eliminated since distribution patterns are difficult to determine from a single occurrence. The chi-square contingency test was used to determine whether a species was randomly distributed or confined to a particular zone and statistically supported the elimination of the

species mentioned above (Ostle and Mensing 1975).

62. Three similar indices were initially investigated: Jaccard's (1901, 1912, 1928) community coefficient, percent similarity (Bray and Curtis 1957), and beta diversity as defined by Bratton (1975). Each index reflected the same general trends, with none emerging as clearly superior. However, Williams and Lambert (1959) suggest that for the recognition of associations, similarity in species combinations (i.e., presence or absence of particular species) is often considered more important than the quantitative contribution of each species. For this reason, Jaccard's community coefficient was chosen and is used in the data analysis for Phases I and II. This particular community coefficient along with its quantitative modifications are among the more widely used indices (Mueller-Dombois and Ellenberg 1974).

63. Transects were initially analyzed both singularly and collectively. The use of a composite required some assurance that the community had been adequately sampled since, theoretically, results should be applicable to the entire community. A low SEM would imply that the sample size was adequate to describe the vegetative community. However, cover estimate variability (as measured by standard error of the mean) was consistently high for a given species in a particular quadrat. This is understandable since a species' absence from only one of the five quadrats comprising the composite would almost automatically result in an unacceptably high SEM (an acceptable SEM was defined as being 10 percent or less of the mean); the number of transects required to reduce the SEM to the required level would be unacceptably high due to the man-hours required.

64. Application of composite results in the field also presents certain problems. The community boundaries determined apply to the composite and not to a particular transect. The critical quadrat or quadrats represent an idealized transition zone for the entire community which may not actually exist at any one particular point in the community. Therefore, the location of this critical zone on the ground would necessarily be somewhat arbitrary. The adopted alternative approach was to consider each transect as a discrete unit. Adequately describing the vegetative community is no longer a concern since each sampling unit is in effect a complete census of the area being analyzed. The boundaries determined by the analysis are a reflection of the biological conditions along a particular transect. The results can be taken and applied to the appropriate transect in the field.

65. A community's transition zone is determined by connecting the critical quadrats for each transect, thus establishing a critical line or zone. (Figure 3). The location of the transition zone will be most accurate at the points where the community was sampled (along the transects). The accuracy of the zone location between transects will be dependent on the uniformity of the zone in this area. The more homogeneous the area is between wetland and upland, the greater the expected accuracy.




Figure 3. Field application of sampling results

66. The analysis of single transects results in certain irregularities which were partially smoothed out by combining quadrats 1 and 2, 3 and 4, etc. This increased the quadrat size from 1- by 4-m to 2- by 4-m. However, Phase II sampling utilized 1- by 4-m quadrats to retain the flexibility of returning to the smaller quadrats, if desirable.

67. It is difficult to determine the degree of similarity at which two samples should be considered from the same association. Jaccard's community coefficient (based on presence-absence data) rarely exceeds 50 or 60 percent. Neighboring communities within the same association often have less than

two-thirds of the species in common (Mueller-Dombois and Ellenberg 1974). Mueller-Dombois and Ellenberg (1974) state that from their experience, a Jaccard community coefficient of more than 25 percent but less than 50 percent (presence-absence data) indicates that samples are from the same association. For values greater than 50 percent they believe the high similarity warrants classification beyond the association level. This was not the purpose of this study and, therefore, only the 25 percent figure was considered critical. Occasionally the Jaccard community coefficient would oscillate around the critical value when moving from the wetland or upland community into the transition zone. These aberrations were judged as representing either microhabitats or a continuation of the existing community, and the transition zone boundary was placed accordingly.

68. Community coefficients were calculated comparing both end quadrats, representing the wetland and upland communities, to all other quadrats. The transition zone boundary was defined as the quadrat or space between quadrats where the community coefficient becomes less than or equal to 25 percent. As a means of supporting the boundaries defined by this comparison, quadrat-to-quadrat coefficients were also calculated and the results compared. The critical point defined by the end quadrat comparison coincided with the lowest or second lowest quadrat-to-quadrat community coefficient in 75 percent of the cases. The close correlation of the two analytical approaches supports the contention that the transition zones identified are real and not mere artifacts of the methodology. Transition zone widths varied between transects within a particular community. Average slopes and transition zone widths for each community are presented in Table 10. The Jaccard community coefficients for all communities sampled during Phase I are presented in Appendix A.

69. There appears to be no correlation between transition zone width and slope (Table 10). The freshwater marsh had the smallest average slope and the second widest transition zones, while the open water community, which had the steepest slope, had a zone of intermediate width. One might have suspected that as the severity of the environmental gradient increased (i.e., slope), the width of the transition zone would decrease. However, this is not reflected by the data.

70. Three different transition zone forms were observed: (a) boundary a discrete line (Figure 4A), (b) boundary with no overlap (Figure 4B), and (c) boundary with overlap (Figure 4C). A line transition zone occurred when

Table 10
Community Slope and Transition Zone Characteristics

| Community | Average Slope m rise/m run | Average Transition Zone Width, m | Transition Zone Width Variability |
|----------------------|-------------------------------|-------------------------------------|--------------------------------------|
| | | | m min - max |
| Saltwater Marsh | 0.06 | 3.0 | 0 - 6 |
| Steep Swamp Forest | 0.17 | 1.6 | 0 - 4 |
| Gradual Swamp Forest | 0.08 | 9.4 | 1 - 18 |
| Open Water | 0.29 | 1.0 | 0 - 4 |
| Shrub | 0.03 | 0.2 | 0 - 1 |
| Freshwater Marsh | 0.02 | 3.4 | 1 - 7 |




Figure 4. Three different configurations of the transition zone

Jaccard's community coefficient reached a critical value in the same quadrat for the quadrat-to-end quadrat comparison. This situation indicated an abrupt transition between upland and wetland communities, with no real transition zone being present. The shrub community provides a particularly good example of this transition zone. Transects 2, 3, 4, and 5 of the gradual swamp forest (Table A6) illustrate transition zones with boundaries that do not overlap. These zones are characterized by having few plant species in common with either the wetland or upland areas and should be viewed as separate communities with their own characteristic plant species. This was the transition zone type found most often in the communities sampled. The last situation involved transition zone boundaries which overlapped, and was observed infrequently. Transect 5 in the saltwater marsh community is an example of this situation. These zones have species in common with both wetland and upland areas and represent a "transition" between the two communities. Both transition zones B and C are an integral part of the wetland ecosystem. Therefore, the critical point would be the quadrat marking the upland edge of this zone.

71. Phase II data analysis revealed the same general patterns as Phase I. Transition zones again assumed one of the three forms previously described. Critical areas were wider for all communities in Phase II with the exception of the gradual swamp forest. The differences varied greatly, with some communities having similar average widths for both phases (saltwater marsh, steep swamp forest, and open water), while others changed by as much as a factor of 12. The patterns were similar, with the shrub and open water communities lowest in both phases. The widest transition zone in Phase II was located in the freshwater marsh, while in Phase I this community exhibited the second widest zone. In Phase I the gradual swamp forest was widest while having an intermediate width in Phase II. The saltwater marsh transition zone had an intermediate width for both. A summary of Phase II sampling results is presented in Table 11.

72. Transition zone widths vary greatly both within and between communities. This fact is a reflection of the highly variable nature of the transition zone in the field.

73. This method of determining transition zone boundaries allows the use of either composite transect analysis or analysis of individual transects. Composite transect analysis results in a line or zone of constant width, while analysis of individual transects results in a zone of varying widths.

Table 11
Minimum, Maximum, and Average Transition Zone Widths

| Community | Transition Zone Width, m (min - max) | Average Transition Zone Width, m |
|------------------|---|-------------------------------------|
| Shrub (A) | 1 - 8 | 4.0 |
| Shrub (B) | 0 - 6 | 2.4 |
| Freshwater Marsh | 7 - 16 | 12.6 |
| Salt Flat (A) | 8 - 14 | 10.8 |
| Salt Flat (B) | 0 - 11 | 4.4 |
| Swamp Forest (A) | 0 - 7 | 3.0 |
| Swamp Forest (B) | 0 - 3 | 3.2 |
| Saltwater Marsh | 1 - 9 | 3.6 |
| Open Water (A) | 0 - 3* | 1.4 |
| Open Water (B) | Not Calculated | Not Calculated |

* The transition zone fell somewhere in a three-quadrat area occupied by a road passing through all transects, making a more precise determination impossible.

Individual transect analysis should be used when a high degree of resolution of zone boundaries is required. However, the use of greater numbers of transects will increase the cost of the determination.

74. The two analytical strategies used for transition zone determination were not as highly correlated in Phase II as they were in Phase I. For narrow transition zones (1 to 2 m), the two strategies defined the same boundary in 60 percent of the cases. Wide transition zones (greater than 2 m) agreed 52 percent of the time. In cases of disagreement the difference was normally not greater than 2 m. The complete results for Phase II are presented in Appendix B.

75. An idealized picture of plant distribution was constructed for each community by combining all five transects (Appendix C: Phase I, Appendix D: Phase II). This provides a generalized overview of the plant distribution encountered in each community and graphically illustrates many of the situations discussed in previous sections.

PART IV: CONCLUSIONS

76. In searching for the best method to define transition zone boundaries, six sampling methodologies, three indices of similarity, and four different analytical strategies were examined. The process is graphically presented in Figure 5.

77. Transition zone narrowness and an effort to improve boundary resolution resulted in the early elimination of four sampling methodologies. Contiguous plots improved boundary definition, while the 1- by 4-m quadrat size involved compromising between the resolution desired and the sampling area required. A strategy based on the analysis of individual transects was superior due to the inherent variability observed in transition zone width. This approach also provides the flexibility of establishing the resolution desired on a case-by-case basis. A two-person team could be expected to sample five to ten 25-m transects per day depending on the density of the vegetation. After the sampling area has been established, the suggested approach is as follows:

- a. Randomly locate transects within the sampling area.
- b. Record presence-absence data for plant species in contiguous 1- by 4-m quadrats along each transect.
- c. Calculate Jaccard community coefficients for each transect (quadrat to end quadrat).
- d. Locate critical quadrat(s) along each transect (Jaccard community coefficient less than or equal to 25 percent).
- e. Locate these critical points along the appropriate transects in the field.
- f. Connect the critical points laterally forming the boundaries of the transition zone.

In conclusion, the described "best method" offers the following advantages:

- a. It does not rely on the presence of "indicator" species; rather, the boundaries are ecologically determined by the position of plant species comprising the wetland, upland, and transition zone communities.
- b. It is uncomplicated, technically reproducible, and therefore amenable to the legal interpretation in defining wetland boundaries.
- c. The field and statistical procedures are well documented in the literature.
- d. The "best method" is applicable to a variety of physiognomic types.
- e. Field application is uncomplicated, and a minimum of training is required for proper implementation by field personnel.

ALTERNATIVES CONSIDERED IN ARRIVING AT THE BEST METHOD




Figure 5. Alternatives considered in arriving at the "best method"

REFERENCES

- Applequist, M. B. 1959. "A Study of Soil and Site Factors Affecting the Growth and Development of Swamp Blackgum and Tupelo-gum Stands in Southwestern Georgia," D.F. dissertation, School of Forestry, Duke Univ., Durham.
- Bozeman, J. R., and J. R. Darrell. 1975. "The River Swamp Ecosystem and Related Vegetation: A Study of Georgia's Coastal Area." Georgia Dept. Nat. Resources, Atlanta.
- Bratton, S. P. 1975. "A Comparison of the Beta Diversity Functions of the Overstory and Herbaceous Understory of a Deciduous Forest," Bull. Torrey Bot. Club, 102:55-60.
- Bray, J. R., and J. T. Curtis. 1957. "An Ordination of the Upland Forest Communities of Southern Wisconsin," Ecol. Monogr., 27:325-349.
- Byrd, H. J., et al. 1961. "Soil Survey of McIntosh County, Georgia," U. S. Dept. Agric., Soil Conserv. Serv.
- Canfield, R. 1941. "Application of the Line Interception Method in Sampling Range Vegetation," J. Forestry, 39:388-394.
- Carter, H. S. 1967. "Climatic Summaries of Resort Areas--The Golden Isles of Georgia," U. S. Dept. Commer. Climatography of the United States, No. 21-9-2.
- _____. 1970. "Georgia Tropical Cyclones and Their Effect on the State," U. S. Dept. Commer., ESSA Tech. Mem. EDSTM 14.
- Cox, G. W. 1980. Laboratory Manual of General Ecology, William C. Brown, Co., Dubuque, Iowa.
- Federal Register. 1977 (19 Jul). "Title 33-Navigable Waters; Chapter II-Corps of Engineers, Department of the Army: Regulatory Programs of the Corps of Engineers," Vol 42, No. 138, pp 37, 122-37, 164, U. S. Government Printing Office, Washington, D. C.
- Georgia Department of Natural Resources. 1976. "Inland Land Use Activities and Georgia's Coastal Waters," Georgia Dept. Nat. Resour., Atlanta.
- Gosselink, J. G., E. P. Odum, and R. N. Pope. 1973. "The Value of the Tidal Marsh," Center for Wetland Resour., Louisiana State Univ., Baton Rouge.
- Haines, E. B. 1975. "Nutrient Inputs to the Coastal Zone: the Georgia and South Carolina Shelf," in: Estuarine Research, 1:303-324, Academic Press, New York.
- Hoyt, J. H. 1967. "Barrier Island Formation," Bull. Geol. Soc. Am., 78(9): 1125-1136.
- _____. 1968. "Geology of the Golden Isles and Lower Georgia Coastal Plain," in: D. S. Maney, ed., The Future of the Marshlands and Sea Islands of Georgia, Georgia Natur. Areas Council and Coastal Area Planning and Devel. Comm., Atlanta, Ga., pp 18-32.
- Hoyt, J. H., and J. R. Hails. 1967. "Pleistocene Shoreline Sediments in Coastal Georgia: Deposition and Modification," Science, 155:1541-1543.
- Jaccard, P. 1901. "Etude Comparative de la Distribution Florale Dans Une Portion des Alpes et du Jura," Bull. Soc. Vand. Sc. Nat., 37:547-579.

- Jaccard, P. 1912. "The Distribution of the Flora of the Alpine Zone," New Phytol., 11:37-50.
- _____. 1928. "Die Statistisch-Floristische Methode als Grundlage der Pflanzensoziologie," in: Alberhalden, ed., Handb. Biol. Arbeitsmeth., 11:165-202.
- Johnson, A. S., et al. 1974. "An Ecological Survey of the Coastal Region of Georgia," National Park Service Monograph Series, No. 3, U. S. Government Printing Office.
- Klawitter, R. A. 1962. "Sweet Gum, Swamp Tupelo, and Water Tupelo Sites in a South Carolina Bottomland Forest," D.F. dissertation, School of Forestry, Duke Univ., Durham.
- Mueller-Dombois, D., and H. Ellenberg. 1974. Aims and Methods of Vegetation Ecology, Wiley, New York.
- Ostle, B., and R. W. Mensing. 1975. Statistics in Research, Iowa State Univ. Press, Ames.
- Radford, A. E., H. E. Ahles, and C. R. Bell. 1968. Manual of the Vascular Flora of the Carolinas, University of North Carolina Press, Chapel Hill.
- Spinner, G. P. 1969. "A Plan for the Marine Resources in the Atlantic Coastal Zone," Am. Geogr. Soc., New York, N. Y.
- Teal, J. M., and J. Kanwisher. 1961. "Gas Exchange in a Georgia Salt Marsh," Limnol. Oceanogr., 6(4):400-415.
- Teal, J. M., and M. Teal. 1969. Life and Death of the Salt Marsh, Little, Brown and Co., Toronto.
- U. S. Geological Survey. 1959. "Surface Water Supply of the United States," Part 2-B, Geol. Surv. Water Supply Paper 1624.
- Visher, G. S., and J. D. Howard. 1974. "Dynamic Relationship Between Hydraulics and Sedimentation in the Altamaha Estuary," J. Sedimentary Petrology, 44:502-521.
- Wharton, C. H. 1978. "The Natural Environments of Georgia," Georgia Dept. Nat. Resour., Atlanta.
- Whittaker, R. W. 1975. Communities and Ecosystems, MacMillan, New York.
- Williams, W. T., and J. M. Lambert. 1959. "Multivariate Methods in Plant Ecology: I. Association Analysis in Plant Communities," J. Ecol., 47:83-101.

APPENDIX A: JACCARD COMMUNITY COEFFICIENTS FOR
PHASE I SAMPLING

Table A1
Jaccard Community Coefficients for the Steep Swamp Forest Community

| Transect | | | | | | | | |
|----------|---|------------|------------|-----------|-----------|------------|------------|------------|
| 1 | Quadrats Compared Jaccard Community Coefficient | | 1,2 67 | 2,3 25 | 3,4 33 | 4,5 67 | 5,6 100 | 6,7 100 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 67 | 1,3 0 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 |
| | Quadrats Compared Jaccard Community Coefficient | 7,1 0 | 7,2 0 | 7,3 25 | 7,4 67 | 7,5 100 | 7,6 100 | 7,7 100 |
| 2 | Quadrats Compared Jaccard Community Coefficient | | 1,2 50 | 2,3 20 | 3,4 50 | 4,5 100 | 5,6 29 | 6,7 25 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 50 | 1,3 20 | 1,4 29 | 1,5 29 | 1,6 0 | 1,7 0 |
| | Quadrats Compared Jaccard Community Coefficient | 7,1 0 | 7,2 0 | 7,3 0 | 7,4 0 | 7,5 0 | 7,6 25 | 7,7 100 |
| 3 | Quadrats Compared Jaccard Community Coefficient | | 1,2 100 | 2,3 50 | 3,4 50 | 4,5 50 | 5,6 83 | 6,7 40 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 100 | 1,3 50 | 1,4 20 | 1,5 13 | 1,6 14 | 1,7 0 |
| | Quadrats Compared Jaccard Community Coefficient | 7,1 0 | 7,2 0 | 7,3 33 | 7,4 33 | 7,5 34 | 7,6 40 | 7,7 100 |
| 4 | Quadrats Compared Jaccard Community Coefficient | | 1,2 100 | 2,3 33 | 3,4 75 | 4,5 38 | 5,6 100 | 6,7 57 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 100 | 1,3 33 | 1,4 50 | 1,5 20 | 1,6 20 | 1,7 0 |
| | Quadrats Compared Jaccard Community Coefficient | 7,1 0 | 7,2 0 | 7,3 17 | 7,4 14 | 7,5 57 | 7,6 57 | 7,7 100 |
| 5 | Quadrats Compared Jaccard Community Coefficient | | 1,2 78 | 2,3 50 | 3,4 50 | 4,5 60 | 5,6 29 | 6,7 75 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 78 | 1,3 30 | 1,4 0 | 1,5 22 | 1,6 10 | 1,7 0 |
| | Quadrats Compared Jaccard Community Coefficient | 7,1 0 | 7,2 9 | 7,3 13 | 7,4 20 | 7,5 14 | 7,6 75 | 7,7 100 |

Table A2
Jaccard Community Coefficients for the Salt Shrub Community

| Transect | | | | | | | |
|----------|-------------------------------|-----|-----|-----|-----|-----|-----|
| 1 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 6,7 |
| | Jaccard Community Coefficient | | 67 | 57 | 67 | 80 | 67 |
| | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,7 |
| | Jaccard Community Coefficient | 100 | 67 | 29 | 50 | 60 | 0 |
| | | | | | | | |
| | Quadrats Compared | 7,1 | 7,2 | 7,3 | 7,4 | 7,5 | 7,7 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 100 |
| | | | | | | | |
| 2 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 6,7 |
| | Jaccard Community Coefficient | | 50 | 83 | 50 | 38 | 60 |
| | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,7 |
| | Jaccard Community Coefficient | 100 | 50 | 60 | 60 | 13 | 0 |
| | | | | | | | |
| | Quadrats Compared | 7,1 | 7,2 | 7,3 | 7,4 | 7,5 | 7,7 |
| | Jaccard Community Coefficient | 0 | 13 | 0 | 0 | 29 | 100 |
| | | | | | | | |
| 3 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 6,7 |
| | Jaccard Community Coefficient | | 60 | 60 | 50 | 25 | 67 |
| | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,7 |
| | Jaccard Community Coefficient | 100 | 60 | 100 | 50 | 0 | 0 |
| | | | | | | | |
| | Quadrats Compared | 7,1 | 7,2 | 7,3 | 7,4 | 7,5 | 7,7 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 29 | 100 |
| | | | | | | | |
| 4 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 6,7 |
| | Jaccard Community Coefficient | | 100 | 50 | 57 | 50 | 33 |
| | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,7 |
| | Jaccard Community Coefficient | 100 | 100 | 50 | 43 | 11 | 0 |
| | | | | | | | |
| | Quadrats Compared | 7,1 | 7,2 | 7,3 | 7,4 | 7,5 | 7,7 |
| | Jaccard Community Coefficient | 0 | 0 | 13 | 11 | 25 | 100 |
| | | | | | | | |
| 5 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 6,7 |
| | Jaccard Community Coefficient | | 86 | 78 | 67 | 83 | 33 |
| | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,7 |
| | Jaccard Community Coefficient | 100 | 86 | 67 | 71 | 57 | 0 |
| | | | | | | | |
| | Quadrats Compared | 7,1 | 7,2 | 7,3 | 7,4 | 7,5 | 7,7 |
| | Jaccard Community Coefficient | 0 | 14 | 11 | 0 | 0 | 100 |
| | | | | | | | |

Table A3
Jaccard Community Coefficients for the Freshwater Marsh Community

| <u>Transect</u> | | | | | | | | | | |
|-----------------------------|---|------------|------------|-----------|------------|------------|------------|-------------|------------|-----------------------------|
| 1 | Quadrats Compared Jaccard Community Coefficient | | 1,2 44 | 2,3 57 | 3,4 11 | 4,5 14 | 5,6 60 | 6,7 20 | 7,8 33 | 8,9 25 9,10 33 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 44 | 1,3 33 | 1,4 33 | 1,5 0 | 1,6 9 | 1,7 0 | 1,8 11 | 1,9 0 1,10 0 |
| | Quadrats Compared Jaccard Community Coefficient | 10,1 0 | 10,2 0 | 10,3 0 | 10,4 0 | 10,5 25 | 10,6 40 | 10,7 100 | 10,8 25 | 10,9 33 10,10 100 |
| 2 | Quadrats Compared Jaccard Community Coefficient | | 1,2 82 | 2,3 67 | 3,4 43 | 4,5 57 | 5,6 58 | 6,7 40 | 7,8 67 | 8,9 40 9,10 100 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 82 | 1,3 42 | 1,4 24 | 1,5 12 | 1,6 6 | 1,7 8 | 1,8 0 | 1,9 0 1,10 0 |
| | Quadrats Compared Jaccard Community Coefficient | 10,1 0 | 10,2 0 | 10,3 0 | 10,4 8 | 10,5 20 | 10,6 22 | 10,7 17 | 10,8 40 | 10,9 100 10,10 100 |
| 3 | Quadrats Compared Jaccard Community Coefficient | | 1,2 64 | 2,3 58 | 3,4 64 | 4,5 60 | 5,6 33 | 6,7 50 | 7,8 57 | 8,9 63 9,10 75 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 64 | 1,3 36 | 1,4 31 | 1,5 30 | 1,6 8 | 1,7 0 | 1,8 0 | 1,9 0 1,10 0 |
| | Quadrats Compared Jaccard Community Coefficient | 10,1 0 | 10,2 6 | 10,3 7 | 10,4 6 | 10,5 8 | 10,6 18 | 10,7 50 | 10,8 75 | 10,9 75 10,10 100 |
| 4 | Quadrats Compared Jaccard Community Coefficient | | 1,2 82 | 2,3 67 | 3,4 58 | 4,5 64 | 5,6 50 | 6,7 50 | 7,8 13 | 8,9 50 9,10 13 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 82 | 1,3 50 | 1,4 36 | 1,5 31 | 1,6 23 | 1,7 17 | 1,8 0 | 1,9 0 1,10 0 |
| | Quadrats Compared Jaccard Community Coefficient | 10,1 0 | 10,2 0 | 10,3 0 | 10,4 9 | 10,5 11 | 10,6 13 | 10,7 17 | 10,8 50 | 10,9 33 10,10 100 |
| <u>Transect (Continued)</u> | | | | | | | | | | |
| 5 | Quadrats Compared Jaccard Community Coefficient | | 1,2 47 | 2,3 50 | 3,4 38 | 4,5 67 | 5,6 82 | 6,7 54 | 7,8 55 | 8,9 88 9,10 86 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 47 | 1,3 43 | 1,4 16 | 1,5 20 | 1,6 18 | 1,7 6 | 1,8 0 | 1,9 0 1,10 0 |
| | Quadrats Compared Jaccard Community Coefficient | 10,1 0 | 10,2 13 | 10,3 6 | 10,4 21 | 10,5 25 | 10,6 42 | 10,7 50 | 10,8 75 | 10,9 86 10,10 100 |

Table A4
Jaccard Community Coefficients for the Saltwater Marsh Community

| Transect | | | | | | | | | | | |
|----------|-------------------------------|------|------|------|------|------|------|------|------|------|-------|
| 1 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 25 | 60 | 17 | 33 | 44 | 67 | 60 | 64 | 67 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 25 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 0 | 11 | 29 | 22 | 27 | 44 | 50 | 67 | 100 |
| 2 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 33 | 43 | 30 | 63 | 75 | 30 | 50 | 40 | 60 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 33 | 14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 0 | 10 | 11 | 22 | 65 | 67 | 75 | 60 | 100 |
| 3 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 67 | 67 | 9 | 64 | 75 | 56 | 78 | 38 | 17 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 67 | 33 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 36 | 18 | 9 | 20 | 33 | 17 | 100 |
| 4 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 50 | 33 | 100 | 25 | 60 | 71 | 100 | 50 | 83 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 50 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 14 | 13 | 13 | 38 | 50 | 63 | 56 | 83 | 100 |
| 5 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 60 | 29 | 40 | 50 | 20 | 75 | 60 | 75 | 67 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 60 | 43 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 17 | 50 | 67 | 25 | 67 | 50 | 50 | 67 | 100 |

Table A5
Jaccard Community Coefficients for the Open Water Community

| Transect | | | | | | | | | |
|----------|---|------------|-----------|-----------|-----------|-----------|------------|------------|------------|
| 1 | Quadrats Compared Jaccard Community Coefficient | | 1,2 50 | 2,3 33 | 3,4 44 | 4,5 57 | 5,6 100 | 6,7 25 | 7,8 100 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 50 | 1,3 0 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 |
| | Quadrats Compared Jaccard Community Coefficient | 8,1 0 | 8,2 0 | 8,3 9 | 8,4 30 | 8,5 25 | 8,6 25 | 8,7 100 | 8,8 100 |
| 2 | Quadrats Compared Jaccard Community Coefficient | | 1,2 75 | 2,3 20 | 3,4 17 | 4,5 50 | 5,6 60 | 6,7 60 | 7,8 50 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 75 | 1,3 0 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 |
| | Quadrats Compared Jaccard Community Coefficient | 8,1 0 | 8,2 13 | 8,3 40 | 8,4 67 | 8,5 50 | 8,6 50 | 8,7 50 | 8,8 100 |
| 3 | Quadrats Compared Jaccard Community Coefficient | | 1,2 67 | 2,3 22 | 3,4 40 | 4,5 33 | 5,6 75 | 6,7 56 | 7,8 43 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 67 | 1,3 0 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 |
| | Quadrats Compared Jaccard Community Coefficient | 8,1 0 | 8,2 20 | 8,3 17 | 8,4 20 | 8,5 43 | 8,6 43 | 8,7 43 | 8,8 100 |
| 4 | Quadrats Compared Jaccard Community Coefficient | | 1,2 50 | 2,3 14 | 3,4 57 | 4,5 83 | 5,6 50 | 6,7 67 | 7,8 63 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 50 | 1,3 0 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 |
| | Quadrats Compared Jaccard Community Coefficient | 8,1 0 | 8,2 0 | 8,3 22 | 8,4 43 | 8,5 38 | 8,6 33 | 8,7 63 | 8,8 100 |
| 5 | Quadrats Compared Jaccard Community Coefficient | | 1,2 33 | 2,3 40 | 3,4 88 | 4,5 44 | 5,6 43 | 6,7 0 | 7,8 20 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 33 | 1,3 0 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 |
| | Quadrats Compared Jaccard Community Coefficient | 8,1 0 | 8,2 29 | 8,3 43 | 8,4 33 | 8,5 17 | 8,6 50 | 8,7 20 | 8,8 100 |

Table A6
Jaccard Community Coefficients for the Gradual Swamp Forest Community*

| Transect | | | | | | | | | | | | |
|----------|-------------------------------|------|------|------|------|------|------|------|------|-------|-------|-------|
| 1 | Quadrats Compared | - | - | - | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | - | - |
| | Jaccard Community Coefficient | - | - | - | 50 | 67 | 67 | 67 | 50 | 25 | - | - |
| | Quadrats Compared | - | - | 4,4 | 4,5 | 4,6 | 4,7 | 4,8 | 4,9 | 4,10 | - | - |
| | Jaccard Community Coefficient | - | - | 100 | 50 | 75 | 50 | 75 | 40 | 0 | - | - |
| | Quadrats Compared | - | - | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 | - | - |
| | Jaccard Community Coefficient | - | - | 20 | 0 | 0 | 0 | 0 | 25 | 100 | - | - |
| | Quadrats Compared | - | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | - | - |
| 2 | Jaccard Community Coefficient | - | 0 | 33 | 100 | 100 | 67 | 20 | 50 | 60 | - | - |
| | Quadrats Compared | 2,2 | 2,3 | 2,4 | 2,5 | 2,6 | 2,7 | 2,8 | 2,9 | 2,10 | - | - |
| | Jaccard Community Coefficient | 100 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | - | - |
| | Quadrats Compared | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 | - | - |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 40 | 60 | 100 | - | - |
| 3 | Quadrats Compared | - | - | - | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | - | - |
| | Jaccard Community Coefficient | - | - | - | 25 | 60 | 33 | 67 | 67 | 50 | - | - |
| | Quadrats Compared | - | - | 4,4 | 4,5 | 4,6 | 4,7 | 4,8 | 4,9 | 4,10 | - | - |
| | Jaccard Community Coefficient | - | - | 100 | 25 | 0 | 0 | 0 | 0 | 0 | - | - |
| | Quadrats Compared | - | - | - | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 | - |
| | Jaccard Community Coefficient | - | - | - | 0 | 0 | 20 | 50 | 33 | 50 | 100 | - |
| | Quadrats Compared | - | - | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 |
| 4 | Jaccard Community Coefficient | - | - | 40 | 60 | 75 | 60 | 67 | 67 | 60 | 33 | 50 |
| | Quadrats Compared | - | 2,2 | 2,3 | 2,4 | 2,5 | 2,6 | 2,7 | 2,8 | 2,9 | 2,10 | 2,11 |
| | Jaccard Community Coefficient | - | 100 | 40 | 17 | 20 | 14 | 14 | 0 | 0 | 14 | 17 |
| | Quadrats Compared | - | 12,2 | 12,3 | 12,4 | 12,5 | 12,6 | 12,7 | 12,8 | 12,9 | 12,10 | 12,11 |
| | Jaccard Community Coefficient | - | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 20 | 25 |
| 5 | Quadrats Compared | - | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 |
| | Jaccard Community Coefficient | - | 33 | 20 | 50 | 50 | 50 | 33 | 17 | 80 | 29 | 43 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 |
| | Jaccard Community Coefficient | 100 | 33 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 12,1 | 12,2 | 12,3 | 12,4 | 12,5 | 12,6 | 12,7 | 12,8 | 12,9 | 12,10 | 12,11 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 20 | 20 |
| | Quadrats Compared | 12,1 | 12,2 | 12,3 | 12,4 | 12,5 | 12,6 | 12,7 | 12,8 | 12,9 | 12,10 | 12,11 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 20 | 20 |

* Vegetation patterns made it necessary to adjust end quadrats to which comparisons were made.

APPENDIX B: JACCARD COMMUNITY COEFFICIENTS FOR
PHASE II SAMPLING

Table B1
Jaccard Community Coefficients for the Swamp Forest (A) Community*

| Transect | | | | | | | | | | | |
|----------|---|------------|------------|------------|------------|------------|-------------|-------------|------------|------------|--|
| 1 | Quadrats Compared Jaccard Community Coefficient | 1,2 67 | 2,3 67 | 3,4 25 | 4,5 50 | 5,6 33 | 6,7 33 | 7,8 67 | 8,9 25 | 9,10 40 | 10,11 50 11,12 25 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 67 | 1,3 33 | 1,4 25 | 1,5 25 | 1,6 0 | 1,7 25 | 1,8 0 | 1,9 25 | 1,10 0 1,11 0 1,12 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 0 | 12,3 0 | 12,4 0 | 12,5 20 | 12,6 33 | 12,7 20 | 12,8 25 | 12,9 50 | 12,10 75 12,11 25 12,12 100 |
| 2 | Quadrats Compared Jaccard Community Coefficient | 1,2 67 | 2,3 50 | 3,4 33 | 4,5 0 | 5,6 33 | 6,7 25 | 7,8 50 | 8,9 33 | 9,10 40 | 10,11 80 |
| | Quadrats Compared Jaccard Community Coefficient | 1,2 100 | 1,3 67 | 1,4 25 | 1,5 50 | 1,6 0 | 1,7 25 | 1,8 0 | 1,9 0 | 1,10 0 | 1,11 - - |
| | Quadrats Compared Jaccard Community Coefficient | 11,1 0 | 11,2 0 | 11,3 0 | 11,4 0 | 11,5 25 | 11,6 40 | 11,7 25 | 11,8 20 | 11,9 50 | 11,10 80 11,11 100 |
| 3 | Quadrats Compared Jaccard Community Coefficient | 1,2 100 | 2,3 50 | 3,4 33 | 4,5 100 | 5,6 100 | 6,7 50 | 7,8 67 | 8,9 75 | 9,10 75 | 10,11 67 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 100 | 1,3 50 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 11,1 0 | 11,2 0 | 11,3 0 | 11,4 33 | 11,5 50 | 11,6 100 | 11,7 67 | 11,8 50 | 11,9 67 | 11,10 100 11,11 100 |
| 4 | Quadrats Compared Jaccard Community Coefficient | 1,2 50 | 2,3 25 | 3,4 0 | 4,5 50 | 5,6 25 | 6,7 50 | 7,8 20 | 8,9 40 | 9,10 67 | 10,11 50 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 50 | 1,3 0 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 11,1 0 | 11,2 0 | 11,3 0 | 11,4 33 | 11,5 25 | 11,6 20 | 11,7 20 | 11,8 20 | 11,9 40 | 11,10 50 11,11 100 |
| 5 | Quadrats Compared Jaccard Community Coefficient | | 2,3 100 | 3,4 0 | 4,5 0 | 5,6 100 | 6,7 50 | 7,8 33 | 8,9 60 | 9,10 80 | 10,11 25 |
| | Quadrats Compared Jaccard Community Coefficient | | 2,2 100 | 2,3 100 | 2,4 0 | 2,5 33 | 2,6 33 | 2,7 0 | 2,8 0 | 2,9 0 | 2,10 0 2,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | | 11,2 0 | 11,3 0 | 11,4 0 | 11,5 50 | 11,6 50 | 11,7 100 | 11,8 33 | 11,9 20 | 11,10 25 11,11 100 |

* Vegetation patterns made it necessary to adjust end quadrats to which comparisons were made.

Table B2
Jaccard Community Coefficients for the Freshwater Marsh Community

| Transect | | | | | | | | | | | | | |
|----------|-------------------------------|------|------|------|------|------|------|------|------|------|-------|-------|-------|
| 1 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 | 11,12 |
| | Jaccard Community Coefficient | | 100 | 31 | 54 | 42 | 44 | 44 | 63 | 17 | 0 | 33 | 33 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 | 1,12 |
| | Jaccard Community Coefficient | 100 | 100 | 38 | 21 | 8 | 9 | 8 | 8 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 12,1 | 12,2 | 12,3 | 12,4 | 12,5 | 12,6 | 12,7 | 12,8 | 12,9 | 12,10 | 12,11 | 12,12 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 9 | 13 | 0 | 0 | 0 | 0 | 33 | 33 | 100 |
| 2 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 | 11,12 |
| | Jaccard Community Coefficient | | 50 | 38 | 50 | 40 | 29 | 40 | 40 | 33 | 29 | 17 | 25 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 | 1,12 |
| | Jaccard Community Coefficient | 100 | 50 | 33 | 27 | 20 | 11 | 14 | 11 | 11 | 10 | 0 | 0 |
| | Quadrats Compared | 12,1 | 12,2 | 12,3 | 12,4 | 12,5 | 12,6 | 12,7 | 12,8 | 12,9 | 12,10 | 12,11 | 12,12 |
| | Jaccard Community Coefficient | 0 | 15 | 0 | 22 | 0 | 17 | 0 | 0 | 0 | 17 | 25 | 100 |
| 3 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 | 11,12 |
| | Jaccard Community Coefficient | | 43 | 50 | 75 | 50 | 43 | 33 | 20 | 43 | 33 | 0 | 33 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 | 1,12 |
| | Jaccard Community Coefficient | 100 | 43 | 22 | 13 | 13 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 12,1 | 12,2 | 12,3 | 12,4 | 12,5 | 12,6 | 12,7 | 12,8 | 12,9 | 12,10 | 12,11 | 12,12 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 13 | 0 | 17 | 0 | 33 | 100 |
| 4 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 | 11,12 |
| | Jaccard Community Coefficient | | 75 | 33 | 50 | 33 | 50 | 27 | 14 | 33 | 17 | 0 | 0 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 | 1,12 |
| | Jaccard Community Coefficient | 100 | 75 | 22 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 12,1 | 12,2 | 12,3 | 12,4 | 12,5 | 12,6 | 12,7 | 12,8 | 12,9 | 12,10 | 12,11 | 12,12 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 25 | 0 | 100 |
| 5 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 | 11,12 |
| | Jaccard Community Coefficient | | 75 | 57 | 43 | 45 | 70 | 56 | 25 | 75 | 40 | 75 | 40 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 | 1,12 |
| | Jaccard Community Coefficient | 100 | 75 | 43 | 13 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 12,1 | 12,2 | 12,3 | 12,4 | 12,5 | 12,6 | 12,7 | 12,8 | 12,9 | 12,10 | 12,11 | 12,12 |
| | Jaccard Community Coefficient | 0 | 0 | 11 | 0 | 8 | 11 | 25 | 50 | 40 | 50 | 40 | 100 |

Table B3
Jaccard Community Coefficients for the Shrub (A) Community

| Transect | | | | | | | | | | |
|----------|-------------------------------|------|------|------|------|------|------|------|------|------|
| 1 | Quadrats Compared | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | 100 | 50 | 100 | 25 | 67 | 0 | 60 | 60 | 33 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 |
| 2 | Jaccard Community Coefficient | 100 | 100 | 50 | 50 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 20 | 33 |
| 3 | Quadrats Compared | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | 100 | 67 | 67 | 67 | 50 | 50 | 60 | 60 | 33 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 |
| 4 | Jaccard Community Coefficient | 100 | 100 | 67 | 33 | 25 | 20 | 25 | 17 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 20 | 33 |
| 5 | Quadrats Compared | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | 100 | 40 | 17 | 67 | 50 | 60 | 75 | 60 | 60 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 |
| 6 | Jaccard Community Coefficient | 100 | 100 | 40 | 20 | 25 | 17 | 17 | 20 | 14 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 17 | 40 | 20 | 60 |
| 7 | Quadrats Compared | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | 100 | 25 | 40 | 60 | 50 | 100 | 67 | 67 | 50 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 |
| 8 | Jaccard Community Coefficient | 100 | 100 | 25 | 40 | 50 | 33 | 33 | 25 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 33 | 50 |
| 9 | Quadrats Compared | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | 100 | 100 | 0 | 33 | 67 | 50 | 100 | 100 | 20 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 |
| 10 | Jaccard Community Coefficient | 100 | 100 | 100 | 0 | 25 | 33 | 20 | 20 | 20 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 25 | 0 | 20 | 20 | 20 |

Table B4
Jaccard Community Coefficients for the Shrub (B) Community

| Transect | | | | | | | | | | | | |
|----------|-------------------------------|------|------|------|------|------|------|------|------|------|-------|-------|
| 1 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 |
| | Jaccard Community Coefficient | | 71 | 44 | 78 | 60 | 63 | 67 | 0 | 0 | 67 | 67 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 |
| | Jaccard Community Coefficient | 100 | 71 | 44 | 36 | 30 | 20 | 10 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 11,1 | 11,2 | 11,3 | 11,4 | 11,5 | 11,6 | 11,7 | 11,8 | 11,9 | 11,10 | 11,11 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 14 | 20 | 33 | 33 | 67 | 100 |
| 2* | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | - | - |
| | Jaccard Community Coefficient | | 83 | 63 | 50 | 29 | 25 | 0 | 0 | 50 | - | - |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | - | - |
| | Jaccard Community Coefficient | 100 | 83 | 50 | 42 | 13 | 20 | 20 | 0 | 0 | - | - |
| | Quadrats Compared | 9,1 | 9,2 | 9,3 | 9,4 | 9,5 | 9,6 | 9,7 | 9,8 | 9,9 | - | - |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 50 | 100 | - | - |
| 3 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 |
| | Jaccard Community Coefficient | | 67 | 60 | 40 | 75 | 0 | 50 | 17 | 50 | 33 | 67 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 |
| | Jaccard Community Coefficient | 100 | 67 | 33 | 50 | 33 | 14 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 11,1 | 11,2 | 11,3 | 11,4 | 11,5 | 11,6 | 11,7 | 11,8 | 11,9 | 11,10 | 11,11 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 20 | 20 | 40 | 25 | 67 | 100 |
| 4 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 |
| | Jaccard Community Coefficient | | 67 | 100 | 57 | 50 | 20 | 0 | 33 | 33 | 33 | 100 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 |
| | Jaccard Community Coefficient | 100 | 67 | 67 | 80 | 33 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 11,1 | 11,2 | 11,3 | 11,4 | 11,5 | 11,6 | 11,7 | 11,8 | 11,9 | 11,10 | 11,11 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 0 | 33 | 33 | 33 | 100 | 100 |
| 5 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 | 10,11 |
| | Jaccard Community Coefficient | | 75 | 60 | 42 | 60 | 17 | 20 | 50 | 50 | 25 | 67 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 | 1,11 |
| | Jaccard Community Coefficient | 100 | 75 | 80 | 42 | 17 | 33 | 0 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 11,1 | 11,2 | 11,3 | 11,4 | 11,5 | 11,6 | 11,7 | 11,8 | 11,9 | 11,10 | 11,11 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 0 | 20 | 100 | 50 | 33 | 67 | 100 |

* Vegetation patterns made it necessary to adjust end quadrats to which comparisons were made.

Table B5
Jaccard Community Coefficients for the Swamp Forest (B) Community

| Transect | | | | | | | | | | | |
|----------|-------------------------------|------|------|------|------|------|------|------|------|------|-------|
| 1 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 100 | 20 | 25 | 50 | 33 | 33 | 13 | 50 | 38 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 100 | 20 | 0 | 0 | 20 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 0 | 13 | 33 | 17 | 29 | 57 | 38 | 38 | 100 |
| 2 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 100 | 60 | 75 | 100 | 50 | 40 | 42 | 33 | 44 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 100 | 60 | 40 | 40 | 17 | 0 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 0 | 11 | 13 | 13 | 29 | 42 | 56 | 44 | 100 |
| 3 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 100 | 100 | 100 | 33 | 100 | 50 | 50 | 50 | 40 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 100 | 100 | 100 | 33 | 33 | 33 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 0 | 40 | 40 | 40 | 40 | 40 | 100 |
| 4 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 100 | 50 | 50 | 67 | 67 | 67 | 50 | 100 | 100 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 100 | 50 | 33 | 25 | 33 | 25 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 25 | 50 | 25 | 50 | 100 | 100 | 100 |
| 5 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 80 | 29 | 60 | 100 | 50 | 67 | 33 | 67 | 100 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 80 | 42 | 14 | 14 | 14 | 17 | 0 | 0 | 0 |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 0 | 33 | 50 | 50 | 50 | 25 | 67 | 100 | 100 |

Table B6
Jaccard Community Coefficients for the Salt Flat (A) Community

| Transect | | | | | | | | | | | | |
|----------|---|------------|-----------|-----------|-----------|-----------|-----------|------------|------------|-------------|-------------|--------------|
| 1 | Quadrats Compared Jaccard Community Coefficient | 1,2 40 | 2,3 57 | 3,4 57 | 4,5 60 | 5,6 67 | 6,7 33 | 7,8 50 | 8,9 33 | 9,10 60 | 10,11 40 | 11,12 33 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 40 | 1,3 14 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 0 | 12,3 0 | 12,4 0 | 12,5 0 | 12,6 0 | 12,7 33 | 12,8 0 | 12,9 0 | 12,10 20 | 12,11 33 |
| 2 | Quadrats Compared Jaccard Community Coefficient | 1,2 67 | 2,3 33 | 3,4 67 | 4,5 60 | 5,6 50 | 6,7 50 | 7,8 67 | 8,9 25 | 9,10 100 | 10,11 40 | 11,12 50 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 67 | 1,3 17 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 0 | 12,3 0 | 12,4 0 | 12,5 0 | 12,6 0 | 12,7 25 | 12,8 33 | 12,9 0 | 12,10 0 | 12,11 50 |
| 3 | Quadrats Compared Jaccard Community Coefficient | 1,2 40 | 2,3 57 | 3,4 57 | 4,5 60 | 5,6 67 | 6,7 33 | 7,8 67 | 8,9 33 | 9,10 43 | 10,11 17 | 11,12 25 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 40 | 1,3 14 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 0 | 12,3 0 | 12,4 0 | 12,5 0 | 12,6 0 | 12,7 0 | 12,8 0 | 12,9 14 | 12,10 33 | 12,11 25 |
| 4 | Quadrats Compared Jaccard Community Coefficient | 1,2 33 | 2,3 71 | 3,4 43 | 4,5 75 | 5,6 33 | 6,7 25 | 7,8 40 | 8,9 75 | 9,10 20 | 10,11 33 | 11,12 100 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 33 | 1,3 14 | 1,4 0 | 1,5 0 | 1,6 0 | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 0 | 12,3 0 | 12,4 0 | 12,5 0 | 12,6 0 | 12,7 0 | 12,8 25 | 12,9 20 | 12,10 33 | 12,11 100 |
| 5 | Quadrats Compared Jaccard Community Coefficient | 1,2 60 | 2,3 83 | 3,4 71 | 4,5 83 | 5,6 60 | 6,7 67 | 7,8 100 | 8,9 33 | 9,10 67 | 10,11 40 | 11,12 75 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 60 | 1,3 50 | 1,4 29 | 1,5 14 | 1,6 0 | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 0 | 12,3 0 | 12,4 0 | 12,5 0 | 12,6 0 | 12,7 0 | 12,8 0 | 12,9 0 | 12,10 20 | 12,11 75 |

Table B7
Jaccard Community Coefficients for the Salt Flat (B) Community

| Transect | | | | | | | | | | | |
|----------|-------------------|------|------|------|------|------|------|------|------|------|-------|
| 1 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community | | 33 | 83 | 86 | 57 | 33 | 40 | 20 | 67 | 33 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community | 100 | 33 | 29 | 43 | 40 | 17 | 20 | 20 | 25 | 0 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community | 0 | 0 | 0 | 0 | 0 | 20 | 25 | 25 | 33 | 100 |
| | Coefficient | | | | | | | | | | |
| 2 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community | | 67 | 71 | 50 | 100 | 20 | 60 | 50 | 75 | 50 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community | 100 | 67 | 43 | 17 | 17 | 0 | 13 | 0 | 0 | 0 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community | 0 | 17 | 17 | 33 | 33 | 66 | 40 | 50 | 66 | 100 |
| | Coefficient | | | | | | | | | | |
| 3 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community | | 40 | 83 | 43 | 100 | 22 | 38 | 50 | 67 | 29 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community | 100 | 40 | 33 | 20 | 20 | 13 | 0 | 0 | 0 | 0 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community | 0 | 0 | 0 | 14 | 14 | 22 | 29 | 29 | 29 | 100 |
| | Coefficient | | | | | | | | | | |
| 4 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community | | 71 | 75 | 75 | 88 | 63 | 33 | 20 | 100 | 100 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community | 100 | 71 | 50 | 33 | 30 | 25 | 0 | 0 | 0 | 0 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community | 0 | 0 | 0 | 11 | 10 | 0 | 20 | 100 | 100 | 100 |
| | Coefficient | | | | | | | | | | |
| 5 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8 | 8,9 | 9,10 |
| | Jaccard Community | | 100 | 75 | 33 | 50 | 50 | 100 | 50 | 40 | 0 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community | 100 | 100 | 75 | 25 | 20 | 14 | 14 | 13 | 0 | 0 |
| | Coefficient | | | | | | | | | | |
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 20 | 0 | 100 |
| | Coefficient | | | | | | | | | | |

Table B8

Jaccard Community Coefficients for the Openwater (A) Community

| Transect | | | | | | | |
|----------|-------------------------------|-----|-----|-----|-----|-----|-----|
| 1 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 |
| | Jaccard Community Coefficient | | 40 | 67 | 50 | 42 | 60 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 |
| | Jaccard Community Coefficient | 100 | 40 | 50 | 14 | 0 | 0 |
| | Quadrats Compared | 6,1 | 6,2 | 6,3 | 6,4 | 6,5 | 6,6 |
| | Jaccard Community Coefficient | 0 | 14 | 25 | 42 | 60 | 100 |
| 2 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 |
| | Jaccard Community Coefficient | | 38 | 60 | 56 | 50 | 67 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 |
| | Jaccard Community Coefficient | 100 | 38 | 22 | 13 | 0 | 0 |
| | Quadrats Compared | 6,1 | 6,2 | 6,3 | 6,4 | 6,5 | 6,6 |
| | Jaccard Community Coefficient | 0 | 8 | 20 | 25 | 67 | 100 |
| 3 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 |
| | Jaccard Community Coefficient | | 50 | 13 | 63 | 30 | 50 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 |
| | Jaccard Community Coefficient | 100 | 50 | 13 | 11 | 0 | 0 |
| | Quadrats Compared | 6,1 | 6,2 | 6,3 | 6,4 | 6,5 | 6,6 |
| | Jaccard Community Coefficient | 0 | 20 | 13 | 11 | 50 | 100 |
| 4 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 |
| | Jaccard Community Coefficient | | 14 | 88 | 63 | 33 | 75 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 |
| | Jaccard Community Coefficient | 100 | 14 | 13 | 0 | 0 | 0 |
| | Quadrats Compared | 6,1 | 6,2 | 6,3 | 6,4 | 6,5 | 6,6 |
| | Jaccard Community Coefficient | 0 | 11 | 10 | 14 | 75 | 100 |
| 5 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 |
| | Jaccard Community Coefficient | | 40 | 83 | 71 | 14 | 33 |
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 |
| | Jaccard Community Coefficient | 100 | 40 | 33 | 33 | 0 | 0 |
| | Quadrats Compared | 6,1 | 6,2 | 6,3 | 6,4 | 6,5 | 6,6 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 17 | 33 | 100 |

Table B9

Jaccard Community Coefficients for the Saltwater Marsh CommunityTransect

1 WAS NOT ANALYZED

| | | | | | | | | |
|---|-------------------------------|--|-----|-----|-----|-----|-----|-----|
| 2 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 |
| | Jaccard Community Coefficient | | 100 | 50 | 33 | 13 | 50 | 57 |

| | | | | | | | | |
|--|-------------------------------|-----|-----|-----|-----|-----|-----|-----|
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 |
| | Jaccard Community Coefficient | 100 | 50 | 14 | 0 | 0 | 0 | 0 |

| | | | | | | | | |
|--|-------------------------------|-----|-----|-----|-----|-----|-----|-----|
| | Quadrats Compared | 7,1 | 7,2 | 7,3 | 7,4 | 7,5 | 7,6 | 7,7 |
| | Jaccard Community Coefficient | 0 | 0 | 13 | 50 | 43 | 57 | 100 |

| | | | | | | | | |
|---|-------------------------------|--|-----|-----|-----|-----|-----|-----|
| 3 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 |
| | Jaccard Community Coefficient | | 33 | 40 | 80 | 67 | 20 | 25 |

| | | | | | | | | |
|--|-------------------------------|-----|-----|-----|-----|-----|-----|-----|
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 |
| | Jaccard Community Coefficient | 100 | 33 | 25 | 20 | 20 | 0 | 0 |

| | | | | | | | | |
|--|-------------------------------|-----|-----|-----|-----|-----|-----|-----|
| | Quadrats Compared | 7,1 | 7,2 | 7,3 | 7,4 | 7,5 | 7,6 | 7,7 |
| | Jaccard Community Coefficient | 0 | 17 | 60 | 50 | 80 | 25 | 100 |

| | | | | | | | | |
|---|-------------------------------|--|-----|-----|-----|-----|-----|-----|
| 4 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 |
| | Jaccard Community Coefficient | | 75 | 60 | 80 | 57 | 71 | 67 |

| | | | | | | | | |
|--|-------------------------------|-----|-----|-----|-----|-----|-----|-----|
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 |
| | Jaccard Community Coefficient | 100 | 75 | 40 | 33 | 29 | 29 | 0 |

| | | | | | | | | |
|--|-------------------------------|-----|-----|-----|-----|-----|-----|-----|
| | Quadrats Compared | 7,1 | 7,2 | 7,3 | 7,4 | 7,5 | 7,6 | 7,7 |
| | Jaccard Community Coefficient | 0 | 0 | 14 | 29 | 43 | 67 | 100 |

| | | | | | | | | | | | |
|---|-------------------------------|--|-----|-----|-----|-----|-----|-----|------|-----|------|
| 5 | Quadrats Compared | | 1,2 | 2,3 | 3,4 | 4,5 | 5,6 | 6,7 | 7,8* | 8,9 | 9,10 |
| | Jaccard Community Coefficient | | 100 | 67 | 33 | 43 | 43 | 71 | 50 | 43 | 60 |

| | | | | | | | | | | | |
|--|-------------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|
| | Quadrats Compared | 1,1 | 1,2 | 1,3 | 1,4 | 1,5 | 1,6 | 1,7 | 1,8 | 1,9 | 1,10 |
| | Jaccard Community Coefficient | 100 | 100 | 67 | 50 | 17 | 50 | 50 | 50 | 17 | 0 |

| | | | | | | | | | | | |
|--|-------------------------------|------|------|------|------|------|------|------|------|------|-------|
| | Quadrats Compared | 10,1 | 10,2 | 10,3 | 10,4 | 10,5 | 10,6 | 10,7 | 10,8 | 10,9 | 10,10 |
| | Jaccard Community Coefficient | 0 | 0 | 0 | 25 | 33 | 11 | 25 | 29 | 60 | 100 |

* Vegetation patterns made it necessary to extend the transect.

Table B10
Jaccard Community Coefficients for the Openwater (B) Community

| Transect | | | | | | | | | | | | |
|----------|---|------------|------------|------------|-----------|--------------|--------------|------------|-------------|------------|--------------|--------------|
| 1 | Quadrats Compared Jaccard Community Coefficient | 1,2 17 | 2,3 50 | 3,4 67 | 4,5 | 5,6 ROAD | 6,7 | 7,8 | 8,9 57 | 9,10 42 | 10,11 83 | 11,12 80 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 0 | 1,3 33 | 1,4 50 | 1,5 | 1,6 ROAD | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 17 | 12,3 0 | 12,4 0 | 12,5 | 12,6 ROAD | 12,7 57 | 12,8 60 | 12,9 67 | 12,10 80 | 12,11 100 |
| 2 | Quadrats Compared Jaccard Community Coefficient | 1,2 33 | 2,3 75 | 3,4 50 | 4,5 50 | 5,6 | 6,7 ROAD | 7,8 50 | 8,9 57 | 9,10 44 | 10,11 75 | 11,12 75 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 33 | 1,3 25 | 1,4 50 | 1,5 100 | 1,6 ROAD | 1,7 0 | 1,8 17 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 13 | 12,3 11 | 12,4 0 | 12,5 0 | 12,6 ROAD | 12,7 13 | 12,8 20 | 12,9 22 | 12,10 75 | 12,11 100 |
| 3 | Quadrats Compared Jaccard Community Coefficient | 1,2 25 | 2,3 50 | 3,4 33 | 4,5 25 | 5,6 | 6,7 ROAD | 7,8 | 8,9 | 9,10 67 | 10,11 80 | 11,12 80 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 25 | 1,3 20 | 1,4 33 | 1,5 0 | 1,6 ROAD | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 13 | 12,3 11 | 12,4 0 | 12,5 0 | 12,6 ROAD | 12,7 67 | 12,8 100 | 12,9 80 | 12,10 100 | 12,11 100 |
| 4 | Quadrats Compared Jaccard Community Coefficient | 1,2 50 | 2,3 67 | 3,4 67 | 4,5 | 5,6 ROAD | 6,7 | 7,8 40 | 8,9 60 | 9,10 50 | 10,11 60 | 11,12 50 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 50 | 1,3 33 | 1,4 50 | 1,5 ROAD | 1,6 0 | 1,7 0 | 1,8 25 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 0 | 12,3 0 | 12,4 0 | 12,5 ROAD | 12,6 60 | 12,7 50 | 12,8 33 | 12,9 43 | 12,10 50 | 12,11 100 |
| 5 | Quadrats Compared Jaccard Community Coefficient | 1,2 33 | 2,3 67 | 3,4 60 | 4,5 | 5,6 ROAD | 6,7 | 7,8 42 | 8,9 71 | 9,10 38 | 10,11 22 | 11,12 57 |
| | Quadrats Compared Jaccard Community Coefficient | 1,1 100 | 1,2 33 | 1,3 33 | 1,4 20 | 1,5 ROAD | 1,6 0 | 1,7 0 | 1,8 0 | 1,9 0 | 1,10 0 | 1,11 0 |
| | Quadrats Compared Jaccard Community Coefficient | 12,1 0 | 12,2 0 | 12,3 0 | 12,4 0 | 12,5 ROAD | 12,6 29 | 12,7 22 | 12,8 38 | 12,9 42 | 12,10 57 | 12,11 100 |

APPENDIX C: SPECIES DISTRIBUTION BY COMMUNITY FOR
PHASE I SAMPLING

Figure C1
Species Distribution in the Saltwater Marsh Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | | | | Dry |
|-----------------------------|----------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|-----|
| | Wet | | | | | | | | | | | | | | | | | | | | |
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | |
| <i>Juncus roemerianus</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Pluchea camphorata</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Nyssa sylvatica</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Ilex vomitoria</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Scleria triglomerata</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Vaccinium corymbosum</i> | | | | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure C2
Species Distribution in the Freshwater Marsh Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | | | | Dry |
|----------------------------------|----------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|--|-----|
| | Wet | | | | | | | | | | | | | | | | | | | | 20 |
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | | |
| <i>Juncus effusus</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Eleocharis fallax</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Scirpus</i> sp. | | | | | | | | | | | | | | | | | | | | | |
| <i>Pontederia cordata</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Aneilema keisak</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Cephalanthus occidentalis</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Hydrocotyle verticillata</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Peltandra virginica</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Cicuta maculata</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Rhus radicans</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Osmunda cinnamomea</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Woodwardia virginica</i> | | | | | | | | | | | | | | | | | | | | | |
| <i>Rubus</i> sp. | | | | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure C3
Species Distribution in the Shrub Community

| Species | Quadrat* | | | | | | | | | | | | | | |
|------------------------------------|----------|---|---|---|---|---|---|---|---|----|----|----|----|----|-----|
| | Wet | | | | | | | | | | | | | | Dry |
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 |
| <i>Borrichia frutescens</i> | | | | | | | | | | | | | | | |
| <i>Baccharis halimifolia</i> | | | | | | | | | | | | | | | |
| <i>Juncus roemerianus</i> | | | | | | | | | | | | | | | |
| <i>Fimbristylis spadicea</i> | | | | | | | | | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | | | | | | | | | |
| <i>Ilex vomitoria</i> | | | | | | | | | | | | | | | |
| <i>Parthenocissus quinquefolia</i> | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure C4
Species Distribution in the Openwater Community

| Species | Quadrat* | | | | | | | |
|---------------------------------|----------|---|---|---|---|---|---|-----|
| | Wet | | | | | | | Dry |
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| <i>Panicum hemitomon</i> | | | | | | | | |
| <i>Hydrochloa caroliniensis</i> | | | | | | | | |
| <i>Lachnanthes caroliniana</i> | | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | | |
| <i>Woodwardia virginica</i> | | | | | | | | |
| <i>Osmunda cinnamomea</i> | | | | | | | | |
| <i>Acer rubrum</i> | | | | | | | | |
| <i>Rubus</i> sp. | | | | | | | | |
| <i>Andropogon glomeratus</i> | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure C5

Species Distribution in the Swamp Forest (Steep) Community

| Species | Quadrat* | | | | | | | | | | | | | | Dry |
|------------------------------|----------|---|---|---|---|---|---|---|---|----|----|----|----|----|-----|
| | Wet | | | | | | | | | | | | | | |
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 |
| <i>Taxodium distichum</i> | | | | | | | | | | | | | | | |
| <i>Nyssa biflora</i> | | | | | | | | | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | | | | | | | | | |
| <i>Osmunda regalis</i> | | | | | | | | | | | | | | | |
| <i>Itea virginica</i> | | | | | | | | | | | | | | | |
| <i>Cyrilla racemiflora</i> | | | | | | | | | | | | | | | |
| <i>Serenoa repens</i> | | | | | | | | | | | | | | | |
| <i>Clethra alnifolia</i> | | | | | | | | | | | | | | | |
| <i>Quercus hemisphaerica</i> | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

APPENDIX D: SPECIES DISTRIBUTION BY COMMUNITY FOR
PHASE II SAMPLING

Figure D1

Species Distribution in the Saltwater Marsh Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | |
|------------------------------|----------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|--------------|
| | Wet 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | Dry 18 19 |
| <i>Juncus roemerianus</i> | | | | | | | | | | | | | | | | | | |
| <i>Ipomoea sagittata</i> | | | | | | | | | | | | | | | | | | |
| <i>Spartina alterniflora</i> | | | | | | | | | | | | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | | | | | | | | | | | | |
| <i>Ilex vomitoria</i> | | | | | | | | | | | | | | | | | | |
| <i>Quercus virginiana</i> | | | | | | | | | | | | | | | | | | |
| <i>Serenoa repens</i> | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure D2

Species Distribution in the Freshwater Marsh (Disturbed) Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | | | | | | | | |
|---------------------------------|----------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----------|
| | Wet 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | Dry 25 |
| <i>Eleocharis quadrangulata</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Cladium jamaicense</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Pontederia cordata</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Nyssa biflora</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Aneilema keisak</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Sabal minor</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Panicum laxiflorum</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Panicum barbulatum</i> | | | | | | | | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure D3
Species Distribution in the Swamp Forest (Steep) Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | | | |
|-----------------------------|----------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|-----------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | Dry 20 |
| <i>Taxodium distichum</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Nyssa biflora</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Cyrilla racemiflora</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Serenoa repens</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Clethra alnifolia</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Vaccinium stamineum</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Vaccinium arboreum</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Osmanthus americana</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Vaccinium corymbosum</i> | | | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure D4
Species Distribution in the Swamp Forest (Gradual) Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | | | | | | | |
|---------------------------|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
| | <div>Wet</div> <div>1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23</div> <div>Dry</div> | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Nyssa ogeche</i> | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Taxodium distichum</i> | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Quercus laurifolia</i> | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Serenoa repens</i> | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Vaccinium arboreum</i> | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Quercus geminata</i> | | | | | | | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure D5
Species Distribution in the Openwater (A) Community

| Species | Quadrat* | | | | | |
|--------------------------------|----------|---|---|---|---|----------|
| | Wet 1 | 2 | 3 | 4 | 5 | Dry 6 |
| <i>*Cyperus</i> spp.** | | | | | | |
| <i>Polygonum punctatum</i> | | | | | | |
| <i>Baccharis halimifolia</i> | | | | | | |
| <i>Euthamia tenuifolia</i> | | | | | | |
| <i>Paspalum urvillei</i> | | | | | | |
| <i>Rubus</i> sp. | | | | | | |
| <i>Panicum scoparium</i> | | | | | | |
| <i>Solidago</i> sp. | | | | | | |
| <i>Ambrosia artemisiifolia</i> | | | | | | |
| <i>Paspalum dilatatum</i> | | | | | | |
| <i>Cynodon dactylon</i> | | | | | | |

* Each quadrat number represents an interval of 1 m.
** Includes both *Cyperus polystachyos* and *Cyperus odoratus*.

Figure D6
Species Distribution in the Openwater (B) Community

| Species | Quadrat* | | | | | | | | |
|--------------------------------|----------|---|---|---|---|---|---|---|----------|
| | Wet 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | Dry 9 |
| <i>Lachnanthes caroliniana</i> | | | | | | | | | |
| <i>Xyris fimbriata</i> | | | | | | | | | |
| <i>Panicum hians</i> | | | | | | | | | |
| <i>Andropogon glomeratus</i> | | | | | | | | | |
| <i>Osmunda cinnamomea</i> | | | | | | | | | |
| <i>Ilex glabra</i> | | | | | | | | | |
| <i>Serenoa repens</i> | | | | | | | | | |
| <i>Persea pubescens</i> | | | | | | | | | |
| <i>Vaccinium stamineum</i> | | | | | | | | | |
| <i>Clethra alnifolia</i> | | | | | | | | | |
| <i>Itea virginica</i> | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure D7
Species Distribution in the Shrub (A) Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | | | Dry | | |
|-----------------------------|----------|--|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|-----|----|----|
| | Wet | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| <i>Borrichia frutescens</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Juncus roemerianus</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Juniperus virginiana</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Pinus taeda</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Ilex vomitoria</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Serenoa repens</i> | | | | | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure D8
Species Distribution in the Shrub (B) Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | | | | | |
|-------------------------------|----------|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|-----|
| | Wet | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | Dry |
| <i>Sporobolus virginicus</i> | | | | | | | | | | | | | | | | | | | | | | 22 |
| <i>Batis maritima</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Iva frutescens</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Borrichia frutescens</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Andropogon glomeratus</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Juncus roemerianus</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Fimbristylis spadiacea</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Myrica cerifera</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Juniperus virginiana</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Ilex vomitoria</i> | | | | | | | | | | | | | | | | | | | | | | |
| <i>Quercus virginiana</i> | | | | | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure D9
Species Distribution in the Salt Flat (A) Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | | | | | | | | |
|------------------------------|----------|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----------|
| | Wet 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | Dry 25 |
| <i>Batis maritima</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Salicornia virginica</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Sporobolus virginicus</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Borrichia frutescens</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Limonium carolinianum</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Juncus roemerianus</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Fimbristylis spadicea</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Spartina alterniflora</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Ilex vomitoria</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Quercus virginiana</i> | | | | | | | | | | | | | | | | | | | | | | | | | |
| <i>Serenoa repens</i> | | | | | | | | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

Figure D10
Species Distribution in the Salt Flat (B) Community

| Species | Quadrat* | | | | | | | | | | | | | | | | | | | |
|--------------------------------|----------|---|---|---|---|---|---|---|---|----|-----|----|----|----|----|----|----|----|----|----|
| | Wet | | | | | | | | | | Dry | | | | | | | | | |
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| <i>Batis maritima</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Spartina alterniflora</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Sporobolus virginicus</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Iva frutescens</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Borrchia frutescens</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Stenotaphrum secundatum</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Fimbristylis spadicea</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Juncus roemerianus</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Ilex vomitoria</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Solidago sempervirens</i> | | | | | | | | | | | | | | | | | | | | |
| <i>Juniperus virginiana</i> | | | | | | | | | | | | | | | | | | | | |

* Each quadrat number represents an interval of 1 m.

APPENDIX E: NATIONAL WETLANDS INVENTORY (NWI) EQUIVALENTS OF
WETLAND TYPES USED IN THIS REPORT

Table E1
National Wetlands Inventory (NWI) Equivalents of Wetland Types
Used in This Report

| <u>Wetland Type</u> | <u>NWI Equivalent</u> |
|---------------------|---|
| Tidal marshes | Estuarine, Intertidal, Emergent, Persistent |
| Salt marshes | Estuarine, Intertidal, Emergent, Persistent |
| Salt flat | Estuarine, Intertidal, Emergent, Persistent |
| Brackish marsh | Estuarine, Intertidal, Emergent, Persistent |
| Freshwater marsh | Palustrine, Emergent, Persistent |
| Hardwood swamp | Palustrine, Forested, Broad-leaf Deciduous |
| River swamp | Palustrine, Forested, Broad-leaf Deciduous |
| Shrub | Palustrine, Scrub-Shrub, Broad-leaf Deciduous |
| Open water | Palustrine, Emergent, Persistent |

END

FILMED

8